

Monday 27 June 2022 - Afternoon

A Level Further Mathematics B (MEI)

Y436/01 Further Pure with Technology

Time allowed: 1 hour 45 minutes

You must have:

- the Printed Answer Booklet
- the Formulae Booklet for Further Mathematics B (MEI)
- a computer with appropriate software
- · a scientific or graphical calculator

INSTRUCTIONS


- Use black ink. You can use an HB pencil, but only for graphs and diagrams.
- Write your answer to each question in the space provided in the Printed Answer
 Booklet. If you need extra space use the lined pages at the end of the Printed Answer
 Booklet. The question numbers must be clearly shown.
- Fill in the boxes on the front of the Printed Answer Booklet.
- Answer all the guestions.
- Where appropriate, your answer should be supported with working. Marks might be given for using a correct method, even if your answer is wrong.
- Give your final answers to a degree of accuracy that is appropriate to the context.
- Do not send this Question Paper for marking. Keep it in the centre or recycle it.

INFORMATION

- The total mark for this paper is 60.
- The marks for each question are shown in brackets [].
- This document has 8 pages.

ADVICE

Read each question carefully before you start your answer.



Answer all the questions.

1 (a) A family of curves is given by the equation

$$x^2 + y^2 + 2axy = 1 (*)$$

where the parameter a is a real number.

You may find it helpful to use a slider (for *a*) to investigate this family of curves.

- (i) On the axes in the Printed Answer Booklet, sketch the curve in each of the cases
 - a = 0
 - *a* = 0.5
 - *a* = 2

[3]

- (ii) State a feature of the curve for the cases a = 0, a = 0.5 that is **not** a feature of the curve in the case a = 2.
- (iii) In the case a=1, the curve consists of two straight lines. Determine the equations of these lines. [2]
- (b) (i) Find an equation of the curve (*) in polar form. [3]
 - (ii) Hence, or otherwise, find, in exact form, the area bounded by the curve, the positive part of the x-axis and the positive part of the y-axis, in the case a = 2. [2]
- (c) In this part of the question m is any real number.

Describing all possible cases, determine the pairs of values a and m for which the curve with equation (*) intersects the straight line given by y = mx. [9]

2 (a) In this part of the question n is an integer greater than 1.

An integer q, where $0 \le q \le n$ is said to be a quadratic residue modulo n if there exists an integer x such that $x^2 \equiv q \pmod{n}$.

Note that under this definition 0 is not considered to be a quadratic residue modulo n.

- (i) Find all the integers x, where $0 \le x < 1000$ which satisfy $x^2 \equiv 481 \pmod{1000}$.
- (ii) Explain why 481 is a quadratic residue modulo 1000. [1]
- (iii) Determine the quadratic residues modulo 11. [2]
- (iv) Determine the quadratic residues modulo 13. [2]
- (v) Show that, for any integer m, $m^2 \equiv (n-m)^2 \pmod{n}$. [2]
- (vi) Prove that if p is prime, where p > 2, then the number of quadratic residues modulo p is $\frac{p-1}{2}$. [4]
- **(b)** Fermat's little theorem states that if p is prime and a is an integer which is co-prime to p, then $a^{p-1} \equiv 1 \pmod{p}$.
 - (i) Use Fermat's little theorem to show that 91 is not prime. [2]
 - (ii) Create a program to find an integer n between 500 and 600 which is not prime and such that $a^{n-1} \equiv 1 \pmod{n}$ for all integers a which are co-prime to n. In the Printed Answer Booklet
 - Write down your program in full.
 - State the integer found by your program.

[6]

3 In this question you are required to consider the family of differential equations

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{y^a}{x+1} - \frac{1}{y} \ (*)$$

and its solutions. The parameter a is a real number.


You should assume that $x \ge 0$ and y > 0 throughout this question.

- (a) In this part of the question a = 1.
 - (i) On the axes in the Printed Answer Booklet
 - Sketch the isocline defined by $\frac{dy}{dx} = 0$.
 - Shade and label the region in which $\frac{dy}{dx} > 0$.
 - Shade and label the region in which $\frac{dy}{dx} < 0$.

For b > 0, find, in terms of b, the solution to (*) which passes through the point (0, b). [1]

- (iii) Determine
 - The values of b > 0 for which the solution in (ii) has a turning point.
 - The corresponding maximum value of *y*.

(b) Fig. 3.1 and Fig. 3.2 show tangent fields for two distinct but unspecified values of a. In each case a sketch of the solution curve y = g(x) which passes through (0, 2) is shown for $0 \le x \le 0.5$.





For the case in **Fig. 3.2** suggest a possible value of *a*. [1]

3

1

0

Fig. 3.2

[3]

[4]

[1]

For the case in **Fig. 3.1** suggest a possible value of a.

- (iii) In each case, continue the sketch of the solution curves for $0.5 \le x \le 5$ in the Printed Answer Booklet. [2]
- (iv) State a feature which is present in one of the curves in part (iii) for $0.5 \le x \le 5$ but not in the other.
- (c) (i) The Euler method for the solution of the differential equation $\frac{dy}{dx} = f(x, y)$ is as follows

$$y_{n+1} = y_n + hf(x_n, y_n).$$

It is given that $x_0 = 0$ and $y_0 = 2$.

- Construct a spreadsheet to solve (*) using the Euler method so that the value of a and the value of b can be varied, in the case $x_0 = 0$ and $y_0 = 2$.
- State the formulae you have used in your spreadsheet.

[3]

(ii) In this part of the question a = 0.1.

Use your spreadsheet with h = 0.1 to approximate the value of y when x = 3 for the solution to (*) in which y = 2 when x = 0.

(iii) In this part of the question a = -0.2.

Use your spreadsheet to approximate, to 1 decimal place, the x-coordinate of the local maximum for the solution to (*) in which y = 2 when x = 0. [3]

END OF QUESTION PAPER

BLANK PAGE

BLANK PAGE



Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of Cambridge University Press & Assessment, which is itself a department of the University of Cambridge.