

Please check the examination details below before entering your candidate information

Candidate surname

Other names

Centre Number

Candidate Number

--	--	--	--	--

--	--	--	--

**Pearson Edexcel Level 3 GCE**

**Tuesday 23 May 2023**

Morning (Time: 1 hour 30 minutes)

Paper  
reference

**8CH0/02**

**Chemistry**

**Advanced Subsidiary**

**PAPER 2: Core Organic and Physical Chemistry**

**You must have:**

Scientific calculator, Data Booklet, ruler

Total Marks

## Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided  
– *there may be more space than you need.*

## Information

- The total mark for this paper is 80.
- The marks for **each** question are shown in brackets  
– *use this as a guide as to how much time to spend on each question.*
- For the question marked with an **asterisk** (\*), marks will be awarded for your ability to structure your answer logically, showing the points that you make are related or follow on from each other where appropriate.
- A Periodic Table is printed on the back cover of this paper.

## Advice

- Read each question carefully before you start to answer it.
- Show all your working in calculations and include units where appropriate.
- Check your answers if you have time at the end.

Turn over ►

P71927A

©2023 Pearson Education Ltd.

N:1/1/1/



  
**Pearson**


Answer ALL questions.

Some questions must be answered with a cross in a box . If you change your mind about an answer, put a line through the box  and then mark your new answer with a cross .

1 This question is about collision theory.

The Maxwell-Boltzmann distribution of molecular energies for a reaction at a particular temperature is shown in the graph.

Add another line on the graph to represent the same reaction with the same number of particles at a **higher** temperature.



(Total for Question 1 = 2 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



2 This question is about alkanes and some of their reactions.

(a) Hexane is a saturated hydrocarbon.

State what is meant by the term 'saturated hydrocarbon'.

Your answer should refer to both words in the term.

(2)

.....

.....

.....



.....

(b) When methane reacts with gaseous chlorine, the initiation step is bond breaking in the chlorine molecule to form chlorine free radicals.

The equation for this step is shown.



Two equations for the propagation step are shown.



(i) Write the overall equation for this reaction of methane with chlorine. State symbols are not required.

(1)

(ii) Calculate the maximum **mass**, in grams, of chloromethane produced when 7.00 g of chlorine gas reacts completely with excess methane.

[Molar mass of  $\text{CH}_3\text{Cl} = 50.5 \text{ g mol}^{-1}$ ]

(2)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 7 1 9 2 7 A 0 3 2 8


- (iii) Write an equation for a termination step for this reaction.  
State symbols are not required.

(1)

- (iv) Further substitution of chloromethane can occur, producing dichloromethane, which is a liquid at room temperature.

Calculate the maximum volume, in  $\text{cm}^3$ , of dichloromethane liquid that could be produced from 12.5 g of chloromethane.  
Give your answer to an appropriate number of significant figures.

(3)



[Density of dichloromethane,  $\text{CH}_2\text{Cl}_2(\text{l}) = 1.32 \text{ g cm}^{-3}$ ]

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA



(c) Commercial diesel fuel is a mixture of hydrocarbons.

One of the components of commercial diesel fuel is a hydrocarbon with the formula  $C_{12}H_{26}$ .

An equation for the complete combustion of  $C_{12}H_{26}$  is shown.



- (i) Calculate the maximum volume, in  $m^3$ , of carbon dioxide gas produced in an engine by the complete combustion of 1.00 **kg** of  $C_{12}H_{26}$  at  $200^\circ C$ , at a pressure of  $6.0 \times 10^6$  Pa.

[The ideal gas equation is  $pV = nRT$   
Gas constant,  $R = 8.31 \text{ J mol}^{-1} \text{ K}^{-1}$ ]

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(ii) Biodiesel is derived from plants and petrodiesel is derived from crude oil. Biodiesel is increasingly widely used, although it is more expensive and provides less energy per kg when burned than petrodiesel.

Give **one** reason why biodiesel is now preferred to petrodiesel.

(1)

.....

.....

.....

.....

**(Total for Question 2 = 14 marks)**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



3 This question is about alkenes and some of their reactions.

(a) Alkenes have the same general formula as cycloalkanes.

Explain why this general formula is different from that of alkanes.  
Include the general formulae in your answer.

(3)

.....

.....

.....

.....

.....

.....

(b) Draw the **skeletal** formula of cyclohexene.


(1)

(c) Draw a diagram of an ethene molecule showing the shape of **all** the bonding electron clouds.  
Include labels for all the bond types within the molecule.

(3)



(d) The reaction scheme shows some of the reactions of but-1-ene.



**A** and **B** are structural isomers. **A** is butan-1-ol.

(i) Draw the fully **displayed** formula of isomer **B**.

(1)

(ii) Draw the **structural** formula of product **C**, which is formed by the reaction of but-1-ene with potassium manganate(VII) in acid conditions.

(1)





(e) The industrial manufacture of margarine involves the addition of hydrogen to an alkene.

Which catalyst is used for this process?

(1)

- A** iron
- B** nickel
- C** phosphoric acid
- D** sodium hydroxide

(f) Polymers can be produced by the addition polymerisation of alkenes. Poly(propene) can be made from propene.

Draw the structure of poly(propene), showing **two** repeat units.

(1)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA



(g) Propene reacts with hydrogen bromide to form a mixture of products.

- (i) Complete the mechanism of the reaction of propene with hydrogen bromide to form 2-bromopropane.  
Include curly arrows and any relevant lone pairs and dipoles.

(4)



- (ii) What is the type and mechanism for the reaction of hydrogen bromide with propene?

(1)

- A** electrophilic addition
- B** electrophilic substitution
- C** nucleophilic addition
- D** nucleophilic substitution





4 This question is about halogenoalkanes and some of their reactions.

(a) **X**, **Y** and **Z** are three different halogenoalkanes.

**X** is 1-chloropropane

**Y** is 1-bromopropane

**Z** is 1-iodopropane

An experiment is carried out to compare the rates of hydrolysis of these compounds.

Outline procedure:

1 cm<sup>3</sup> of each of the three halogenoalkanes, **X**, **Y** and **Z**, is added to separate test tubes, each containing 5 cm<sup>3</sup> of ethanol and 5 cm<sup>3</sup> of aqueous silver nitrate solution, in a water bath at 50 °C.

The time taken for a precipitate to form in each test tube is measured.

(i) Give **three** reasons why these reaction **conditions** are specified.

(3)

(ii) Explain why a precipitate forms.

(2)



(iii) The three halogenoalkanes were placed in order of **increasing** rate of reaction.

Which is the correct sequence?

(1)

- A X, Y, Z
- B X, Z, Y
- C Y, X, Z
- D Z, Y, X

(b) The results table shows the time taken to produce a precipitate when three bromoalkanes react with aqueous ethanolic silver nitrate solution.

Halogenoalkane	Time to produce a precipitate / s
1-bromobutane	58
2-bromobutane	33
2-bromo-2-methylpropane	2

Give a reason why the times taken to produce a precipitate for these isomeric bromoalkanes are different.

(1)

.....

.....

.....

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





DO NOT WRITE IN THIS AREA



P 7 1 9 2 7 A 0 1 3 2 8

(c) (i) Which equation shows the first step of the mechanism for the reaction of 1-bromopropane with ammonia?

(1)




(ii) Write the formula of the final inorganic product of the reaction of 1-bromopropane with **excess** ammonia.

(1)



(iii) The infrared spectrum of ethylamine is shown.



Which absorption peak confirms this as an amine?

(1)

- A absorption P
- B absorption Q
- C absorption R
- D absorption S

(d) 1-bromopropane undergoes reactions when heated with different reagents.

- (i) Give **two** reasons why organic reactions are often heated for a long time but the yield is frequently low.

(2)

.....

.....

.....

.....



P 7 1 9 2 7 A 0 1 5 2 8

(ii) Give the name or formula of the reagent and the condition, other than heat, used to increase the carbon chain length by one carbon atom, starting from 1-bromopropane.

(2)

(iii) 1-bromopropane,  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{Br}$ , can be converted into but-1-ene,  $\text{CH}_3\text{CH}_2\text{CH}=\text{CH}_2$ .

Give the name or formula of the reagent and the condition, other than heat, used for this reaction.

(2)

**(Total for Question 4 = 16 marks)**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**BLANK PAGE**




P 7 1 9 2 7 A 0 1 7 2 8

- 5 A student found a bottle of a colourless liquid that was thought to be an alcohol. However, the label was missing and the identity of the alcohol was unknown.

The student decided to attempt to identify this alcohol (**Q**) by measuring its enthalpy change of combustion, and comparing the result with values in a data book.

The student used the equipment shown in the diagram to determine the enthalpy change of combustion of alcohol **Q**.



### Data

Mass of spirit burner + alcohol **Q** before combustion = 20.24 g

Mass of spirit burner + alcohol **Q** after combustion = 19.48 g

Mass of water in the beaker = 500 g

Temperature of the water before the experiment = 17.8 °C

Temperature of the water at the end of the experiment = 28.7 °C

Specific heat capacity of water = 4.18 Jg<sup>-1</sup>°C<sup>-1</sup>



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(a) (i) Calculate the enthalpy change, in  $\text{kJ g}^{-1}$ , when 1.00 g of alcohol **Q** is burned. Include a sign in your final answer.

(3)

(ii) At the end of the experiment there was a black deposit of carbon on the bottom of the beaker.

Explain the effect of formation of the carbon deposit on your answer to (a)(i).

(2)

.....

.....

.....

.....

.....

.....

.....

.....



(b) Give one theoretical and one practical reason why this procedure is insufficient to identify the alcohol.

(2)

Theoretical reason

.....

.....

.....

Practical reason

.....

.....

.....

(c) The student concluded that alcohol **Q** was either propan-1-ol or propan-2-ol because the mass spectrum had the molecular ion peak at  $m/z = 60$ .

Explain **one** peak that you would expect to be present in the mass spectrum of propan-1-ol but not in the mass spectrum of propan-2-ol.

(2)

.....

.....

.....

.....

.....

.....


.....

**(Total for Question 5 = 9 marks)**



- 6 Ammonium carbamate has the formula  $\text{NH}_2\text{COONH}_4$ . When a sample of ammonium carbamate is placed in a sealed tube and heated, it partially decomposes to the compounds used to synthesise it.

The equation for this reversible reaction is



Which is the expression for the equilibrium constant,  $K_c$ , for this reaction?  
All concentration terms are at equilibrium.

- A  $\frac{[\text{NH}_3][\text{CO}_2]}{[\text{NH}_2\text{COONH}_4]}$
- B  $\frac{[2\text{NH}_3][\text{CO}_2]}{[\text{NH}_2\text{COONH}_4]}$
- C  $[\text{NH}_3]^2[\text{CO}_2]$
- D  $\frac{[\text{NH}_3]^2[\text{CO}_2]}{[\text{NH}_2\text{COONH}_4]}$

(Total for Question 6 = 1 mark)




7 This question is about the primary alcohol butan-1-ol.

(a) Which are the intermolecular force(s) between butan-1-ol molecules in the liquid phase?

(1)


- A hydrogen bonding only
- B hydrogen bonding and permanent dipole-dipole forces only
- C hydrogen bonding, permanent dipole-dipole forces and London forces
- D London forces only

(b) The equation for the complete combustion of butan-1-ol is shown.



Draw a labelled enthalpy level diagram for this reaction, using the axes provided. You may use the labels 'reactants' and 'products' in place of the formulae shown in the equation.

(2)



(c) Butan-1-ol can be oxidised to produce butanal or butanoic acid.

(i) Which is a suitable test and result for butan-1-ol?

(1)

	Test reagent	Observation
<input type="checkbox"/> A	phosphorus(V) chloride $\text{PCl}_5(\text{s})$	white smoke
<input type="checkbox"/> B	phosphorus(V) chloride $\text{PCl}_5(\text{s})$	steamy fumes
<input type="checkbox"/> C	sodium carbonate $\text{Na}_2\text{CO}_3(\text{aq})$	carbon dioxide given off
<input type="checkbox"/> D	sodium carbonate $\text{Na}_2\text{CO}_3(\text{aq})$	effervescence

(ii) State how to produce butanal in high yield when butan-1-ol reacts with acidified potassium dichromate(VI).

(1)

(iii) Describe a chemical test, including the expected results, that would confirm the presence of an aldehyde such as butanal.

(2)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 7 1 9 2 7 A 0 2 3 2 8

\*(iv) Describe, including practical details, how a sample of butanoic acid (boiling temperature 166 °C) could be prepared from butan-1-ol (boiling temperature 117 °C) using acidified potassium dichromate(VI).  
Include labelled diagrams of the apparatus you would use for the reaction, and for collecting the product.  
You may assume that all necessary safety precautions are observed.

(6)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....





DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA

(Total for Question 7 = 13 marks)




P 7 1 9 2 7 A 0 2 5 2 8

8 A data book gave the following information.




(a) State the **two** conditions denoted by the standard symbol  $\ominus$  for this reaction. (2)

(b) (i) Draw a Hess's Law cycle that would enable you to use the data provided to calculate the enthalpy change of formation of gaseous water,  $\Delta_f H$ . (2)



Data:




DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(ii) Calculate the enthalpy change for the reaction shown.



(3)

---

(Total for Question 8 = 7 marks)

---

**TOTAL FOR PAPER = 80 MARKS**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



# The Periodic Table of Elements

1 2 3 4 5 6 7 0 (8) (18)

1.0	<b>H</b>
	hydrogen
	1

### Key

relative atomic mass
<b>atomic symbol</b>
name
atomic (proton) number

(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)
6.9 <b>Li</b> lithium 3	9.0 <b>Be</b> beryllium 4	45.0 <b>Sc</b> scandium 21	47.9 <b>Ti</b> titanium 22	50.9 <b>V</b> vanadium 23	52.0 <b>Cr</b> chromium 24	54.9 <b>Mn</b> manganese 25	55.8 <b>Fe</b> iron 26	58.9 <b>Co</b> cobalt 27	58.7 <b>Ni</b> nickel 28	63.5 <b>Cu</b> copper 29	65.4 <b>Zn</b> zinc 30	10.8 <b>B</b> boron 5	12.0 <b>C</b> carbon 6	14.0 <b>N</b> nitrogen 7	16.0 <b>O</b> oxygen 8	19.0 <b>F</b> fluorine 9	4.0 <b>He</b> helium 2
23.0 <b>Na</b> sodium 11	24.3 <b>Mg</b> magnesium 12	88.9 <b>Y</b> yttrium 39	91.2 <b>Zr</b> zirconium 40	92.9 <b>Nb</b> niobium 41	95.9 <b>Mo</b> molybdenum 42	[98] <b>Tc</b> technetium 43	101.1 <b>Ru</b> ruthenium 44	102.9 <b>Rh</b> rhodium 45	106.4 <b>Pd</b> palladium 46	107.9 <b>Ag</b> silver 47	112.4 <b>Cd</b> cadmium 48	27.0 <b>Al</b> aluminium 13	28.1 <b>Si</b> silicon 14	31.0 <b>P</b> phosphorus 15	32.1 <b>S</b> sulfur 16	35.5 <b>Cl</b> chlorine 17	39.9 <b>Ar</b> argon 18
39.1 <b>K</b> potassium 19	40.1 <b>Ca</b> calcium 20	88.9 <b>Y</b> yttrium 39	91.2 <b>Zr</b> zirconium 40	92.9 <b>Nb</b> niobium 41	95.9 <b>Mo</b> molybdenum 42	[98] <b>Tc</b> technetium 43	101.1 <b>Ru</b> ruthenium 44	102.9 <b>Rh</b> rhodium 45	106.4 <b>Pd</b> palladium 46	107.9 <b>Ag</b> silver 47	112.4 <b>Cd</b> cadmium 48	69.7 <b>Ga</b> gallium 31	72.6 <b>Ge</b> germanium 32	74.9 <b>As</b> arsenic 33	79.0 <b>Se</b> selenium 34	79.9 <b>Br</b> bromine 35	83.8 <b>Kr</b> krypton 36
85.5 <b>Rb</b> rubidium 37	87.6 <b>Sr</b> strontium 38	88.9 <b>Y</b> yttrium 39	91.2 <b>Zr</b> zirconium 40	92.9 <b>Nb</b> niobium 41	95.9 <b>Mo</b> molybdenum 42	[98] <b>Tc</b> technetium 43	101.1 <b>Ru</b> ruthenium 44	102.9 <b>Rh</b> rhodium 45	106.4 <b>Pd</b> palladium 46	107.9 <b>Ag</b> silver 47	112.4 <b>Cd</b> cadmium 48	114.8 <b>In</b> indium 49	118.7 <b>Sn</b> tin 50	121.8 <b>Sb</b> antimony 51	127.6 <b>Te</b> tellurium 52	126.9 <b>I</b> iodine 53	131.3 <b>Xe</b> xenon 54
132.9 <b>Cs</b> caesium 55	137.3 <b>Ba</b> barium 56	138.9 <b>La*</b> lanthanum 57	178.5 <b>Hf</b> hafnium 72	180.9 <b>Ta</b> tantalum 73	183.8 <b>W</b> tungsten 74	186.2 <b>Re</b> rhenium 75	190.2 <b>Os</b> osmium 76	192.2 <b>Ir</b> iridium 77	195.1 <b>Pt</b> platinum 78	197.0 <b>Au</b> gold 79	200.6 <b>Hg</b> mercury 80	204.4 <b>Tl</b> thallium 81	207.2 <b>Pb</b> lead 82	209.0 <b>Bi</b> bismuth 83	[209] <b>Po</b> polonium 84	[210] <b>At</b> astatine 85	[222] <b>Rn</b> radon 86
[223] <b>Fr</b> francium 87	[226] <b>Ra</b> radium 88	[227] <b>Ac*</b> actinium 89	[261] <b>Rf</b> rutherfordium 104	[262] <b>Db</b> dubnium 105	[266] <b>Sg</b> seaborgium 106	[264] <b>Bh</b> bohrium 107	[277] <b>Hs</b> hassium 108	[268] <b>Mt</b> meitnerium 109	[271] <b>Ds</b> darmstadtium 110	[272] <b>Rg</b> roentgenium 111	Elements with atomic numbers 112-116 have been reported but not fully authenticated						

\* Lanthanide series

\* Actinide series

140 <b>Ce</b> cerium 58	141 <b>Pr</b> praseodymium 59	144 <b>Nd</b> neodymium 60	150 <b>Sm</b> samarium 62	152 <b>Eu</b> europium 63	157 <b>Gd</b> gadolinium 64	163 <b>Dy</b> dysprosium 66	165 <b>Ho</b> holmium 67	167 <b>Er</b> erbium 68	169 <b>Tm</b> thulium 69	173 <b>Yb</b> ytterbium 70	175 <b>Lu</b> lutetium 71
232 <b>Th</b> thorium 90	[231] <b>Pa</b> protactinium 91	238 <b>U</b> uranium 92	[242] <b>Pu</b> plutonium 94	[243] <b>Am</b> americium 95	[247] <b>Cm</b> curium 96	[251] <b>Cf</b> californium 98	[254] <b>Es</b> einsteinium 99	[253] <b>Fm</b> fermium 100	[256] <b>Md</b> mendelevium 101	[254] <b>No</b> nobelium 102	[257] <b>Lr</b> lawrencium 103

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

