

Mark Scheme (Results)

June 2016

Pearson Edexcel International GCSE Mathematics B (4MB0) Paper 02R



Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please visit our website at <u>www.edexcel.com</u>.

Our website subject pages hold useful resources, support material and live feeds from our subject advisors giving you access to a portal of information. If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

www.edexcel.com/contactus

Pearson: helping people progress, everywhere

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2016 Publications Code 4MB0_02R_1606_MS All the material in this publication is copyright © Pearson Education Ltd 2016

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.
- Types of mark
 - M marks: method marks
 - A marks: accuracy marks
 - B marks: unconditional accuracy marks (independent of M marks)

Abbreviations

- cao correct answer only
- ft follow through
- isw ignore subsequent working
- SC special case
- oe or equivalent (and appropriate)
- \circ dep dependent
- indep independent
- \circ eeoo each error or omission

• No working

If no working is shown then correct answers normally score full marks

If no working is shown then incorrect (even though nearly correct) answers score no marks.

• With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.

If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks.

Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks.

If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.

If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

If there is no answer on the answer line then check the working for an obvious answer.

• Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. Incorrect cancelling of a fraction that would otherwise be correct.

It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect eg algebra.

Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

• Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

International GCSE Maths						
Q	Working	Answer	Mark	Notes		
1 a	$x^2 - 2 \times 3 \times y = 9 - 6y$		3	M1 Can be embedded in a matrix equation (This also applies to the second equation).		
		<i>x</i> = 3		A1		
		<i>x</i> = -3		A1		
b	y-2x=5		3	M1		
		y = 11 (5+2×"3")		A1		
		· · · · · ·		ft		
		y = -1 (5-2×"-3")		A1		
				ft		
				Total 6 marks		

2 a	$r_{\text{Removed}} = \frac{8}{20} \times 6$ $\left(\frac{8}{20} = \frac{r}{6}\right)$ (o.e.)		M1	
	Alt: (Using half the vertex angle)			
	$r = 8 \times \tan(16.7)$			
		$r_{\text{Removed}} = 2.4 \text{ (cm) (awrt)}$	A1	
b	$(V =)\frac{1}{3}\pi \times 6^2 \times 20 - \frac{1}{3}\pi \times "2.4"^2 \times 8 \qquad (oe)$		M1	
		$=706 \text{ or } 705 \text{ (cm}^{3} \text{) (cao)}$	A1	
c	Time (sec) = $\frac{"706"}{54}$		M1	
		Time (sec) = 13 (cao)	A1	
				Total 6 marks

3 a	$\left(\frac{\mathrm{d}y}{\mathrm{d}x}\right) = 6x^2 - 8x$ (1 term correct)		4	M1
	Fully correct			A1
	$6x^2 - 8x = 0$			M1
	0. 0 0			(dep)
		$x = \frac{4}{3} (cc)$		A1
	OR			
	$(4)^2$ (4)			M1
	$6 \times \left(\frac{4}{3}\right)^2 - 8\left(\frac{4}{3}\right)$			(dep)
		0 (cc)		A1
b	calculating			
	$\frac{dy}{dx}(x=1, \operatorname{say}) = -2$ and $\frac{dy}{dx}(x=2, \operatorname{say}) = +8$			M1
	(ie evaluating $\frac{dy}{dx}$ on either side of the stationary			
	point)			
	Note: any value to the left of $4/3$ (must be > 0) any value to the right of $4/3$			
	Noting the change in sign of $\frac{dy}{dx}$ from negative to			M1 (dep)
	positive as $x \uparrow$ through the stationary point			(dop)

OR Calculating $f(4/3) = 2.63$ (or better) (Accept 71/27) Calculating			M1	<pre>any value to the left of 4/3 (must be > 0) any value to the right of 4/3</pre>
$f(x=1, \operatorname{say}) = 3$ and $f(x=2, \operatorname{say}) = 5$				
and noting that both values are greater than			M1	
f(4/3)			(dep)	
OR (not in syllabus)				
$\frac{\mathrm{d}^2 y}{\mathrm{d}x^2} = 12x - 8$			M1	
$\frac{d^2 y}{dx^2}(x=\frac{4}{3}) > 0$			M1	
$\frac{1}{dx^2} \left(x = \frac{1}{3} \right) > 0$			(dep)	
	Minimum	(cc)	A1	
				Total 5 marks

4	а	Vans = 240		1	B1
	b	One of Lorries: $\frac{1}{5} \times (600 - "240")$ OR Cars: $\frac{4}{5} \times (600 - "240")$ seen (oe)		3	M1
			Lorries = 72 OR Cars = 288		A1
			Cars = (600-"240")-"72" OR Lorries = (600-"240")-"288"		A1 ft
	c	Cars sold = "288" + $\frac{1}{9}$ ×"288" (= 320) (oe) and Lorries sold = $\frac{87.5}{100}$ ×"72" (= 63) (oe)		3	M1 Accept 12.5% increase or decrease here (i.e. 63 or 81)
		$\frac{\% \text{ increase} =}{\frac{("320"+"63"+"240")-600}{600}} \times 100 \text{(oe)}$ (3.833)			M1 (dep)

OR		M1	
Increase in cars sold = $\frac{1}{9} \times "288" (= 32)$			
and (decrease) in lorries sold =			
$\frac{12.5}{100} \times "72" (=9)$			
100			
$\frac{"32"-"9"}{600}$ ×100		M1	
600 ~100		(dep)	
	4%	A1	
			Total 7 marks

5	ai	$xy=1+x$ OR $y = \frac{1}{x}+1$ OR OR		4	M1	
		$xy = y + 1$ $x(y-1) = 1$ OR $y-1 = \frac{1}{x}$			M1 (dep)	
		$y = 1 - \frac{1}{x}$ OR $y(x-1) = 1$ $f^{-1} : x \mapsto \frac{1}{x-1}$			A1	For the letter 'y', accept any other letter
	aii		$(x =)1$ OR $x \neq 1$ OR "not 1"		B1 ft	Cand's (a)(i) must be a fraction with a linear denominator and ft is on the cand's denominator
	b	$\frac{2}{x} + 3 = 4 \times "\frac{1}{x-1}"$		5	M1	Correctly removing at least two different denominators in <i>x</i> This M mark is independent of the previous M mark.
		2(x-1) + 3x(x-1) = 4x (oe)	$3x^2 - 5x - 2 \ (= 0)$		M1 A1	

(3x+1)(x-2)		M1	
attempt to factorise their trinomial quadratic			
OR			
fully correct substitution into a correctly			
quoted formula			
ŌR			
Completing the square method as far as:			
$\left(x - \frac{5}{6}\right)^2 = \frac{2}{3} + \left(\frac{5}{6}\right)^2$			
	$-\frac{1}{3}$ (-0.333 awrt),	A1	
	2 `		
			Total 9 marks

6	а		Triangle A drawn and labelled	1	B1	
	b	$ \begin{pmatrix} -1 & 1 \\ 2 & -1 \end{pmatrix} \begin{pmatrix} 1 & 2 & -1 \\ 1 & 3 & 2 \end{pmatrix} $		3	M1	Method can be implied from the resultant 2 x 3 matrix or the triangle itself
			Triangle <i>B</i> is $(0, 1), (1, 1), (3, -4)$ Triangle <i>B</i> drawn and labelled.		A2 (-1eeoo)	
	с	$\left(\begin{array}{ccc} 2 & 1 \\ -1 & -1 \end{array}\right)" \left(\begin{array}{ccc} 0 & 1 & 3 \\ 1 & 1 & -4 \end{array}\right)"$		3	M1	
			Triangle C is $(1, -1)$, $(3, -2)$, $(2, 1)$ Triangle C drawn and labelled		A2 ft (-1eeoo)	The coordinates can be implied from a 2 x 3 matrix seen or from the triangle itself. If the ft is to be applied we must see an attempt at the matrix multiplication
	d		Rotation	3	B1	
			about origin or (0, 0)		B1	
			(anticlockwise) 270° or clockwise 90° or –90°		B1	A combined transformation earns no marks No marks (or penalties) for the appearance of a matrix here.
						Total 9 marks

7	a		0.25 (oe)	B1
			0.7, 0.3 (oe)	B1
			0.6, 0.4 (oe)	B1
	bi	"0.25" × "0.4"		M1
			0.1 1	A1
			$0.1, \frac{1}{10}$	
	bii	One of $0.75 \times "0.3"$ and $"0.25" \times "0.4"$		M1
		OR		
		$0.75 \times "0.7" + "0.25" \times "0.6"$		
		0.75 ×"0.3" + "0.25"× "0.4"		M1 For "0.25" × "0.4" accept
				(dep) "0.1"
		OR		
		1 (0.75 110.711 110.2511 110.611)		
		$1 - (0.75 \times "0.7" + "0.25" \times "0.6")$		
			0.225 13	A1
			$0.325, \frac{13}{40}$	
	с	a probability		M1
	•	"0.325"		
		"0.1"		M1 "0.325"=
		"0.325"		(dep) $0.75 \times "0.7" + "0.25" \times "0.6"$
		0.325	1	$\begin{array}{c} (acp) & 0.75 \times 0.7 + 0.25 \times 0.0 \\ \hline A1 & Condone \ percentages \ as \end{array}$
			awrt 0.31, $\frac{4}{12}$	probabilities.
			13	
				Total 11 marks

8	$t = ab + adx$ OR $\frac{t}{a} = b + dx$ $t - ab = adx (o.e.)$ OR $\frac{t}{a} - b = dx (o.e.)$		M1 M1 (dep)
		$x = \frac{t - ab}{ad}, x = \frac{t}{ad} - \frac{b}{d} ,$ $x = \frac{\left(\frac{t}{a} - b\right)}{d} \qquad (\text{o.e})$	A1 Do not isw Accept also $x = \frac{-t + ab}{-ad}$ and $x = \frac{\left(\frac{t - ab}{a}\right)}{d}$ Do not accept $\frac{t - ab}{\frac{a}{d}}$ (the order of operation must be clear) $x = \dots$ must be on the answer line for the final A mark
			Total 3 marks

9	$BD^2 = 10^2 + 4^2 - 2 \times 10 \times 4 \times \cos(20)$		3	M1
	$BD = \sqrt{\left(10^2 + 4^2 - 2 \times 10 \times 4 \times \cos(20)\right)} \left(=\sqrt{40.8}\right)$			M1 (dep)
		<i>BD</i> = 6.3894 cm → 6.39		A1
`b	In $\triangle ABD$, $\frac{10}{\sin \angle ADB} = \frac{"6.3894"}{\sin 20}$		4	M1
	$Sin \angle ADB$ $Sin 20$			
	$10^2 = 4^2 + 6.389''^2 - 2 \times 4 \times 6.389'' \times \cos \angle ADB$			
	$\angle ADB = \sin^{-1} \left(\frac{10 \times \sin 20}{"6.3894"} \right) \ (\angle ADB = 147.636)$			M1 (dep)
	OR			
	$\therefore \angle ADB = \cos^{-1} \left(\frac{4^2 + "6.389"^2 - 10^2}{2 \times 4 \times "6.389"} \right) (\angle ADB = 147.636)$			
	$\therefore \angle BDC = 180 - "147.636"$			M1 (dep)
		$\therefore \angle BDC = 32.3637 \rightarrow 32.4^{\circ}$		A1
	$\frac{OR}{\frac{4}{\sin \angle ABD}} = \frac{"6.3894"}{\sin 20}$			M1
				M1
	$\sin \angle ABD = \frac{4 \times \sin 20}{"6.3894"}$ ($\angle ABD = 12.3637$)			(dep)
	$\therefore \angle BDC = 20 + "12.3637"$			M1
		$\therefore \angle BDC = 32.3637 \rightarrow 32.4^{\circ}$		(dep) A1
	1	·· / / / / / / / / / / / / / / / /	1	***

c	$\Delta ABC = 18 = \frac{1}{2} \times 10 \times "(4 + CD)" \times \sin 20$	3	M1
	OR		
	$\Delta ABD = \frac{1}{2} \times 4 \times 10 \times \sin 20 \ (=6.8404)$		
	$\therefore \Delta BCD = 18 - "6.8404" (=11.1596)$		
	$\therefore CD = \frac{18}{\frac{1}{2} \times 10 \times \sin 20} - 4 \qquad (oe)$		M1 (dep)
	OR		
	$(::"11.1596" = \frac{1}{2} \times "6.3894" \times CD \times \sin"32.3637")$		
	$CD = \frac{"11.1596"}{\frac{1}{2} \times "6.3894" \times \sin"32.3637"}$		
	$\frac{-\times"6.3894"\times\sin"32.3637"}{2}$		

∴ CD = 6.5186, 6.5257 cm →6.52, 6.53	$CD = 6.5186$ using 6.39 and 32.4° and $CD = 6.5257$ using 6.3894 and 32.3637° If the incorrectobtuse angle hasbeen penalised inpart (b), Condone
	sin(147.636) in this part of the
	question. Total 12 marks

10 ai		$\overrightarrow{AB} = 2\mathbf{b} - \mathbf{a}$	5	B1	
aii		$\overrightarrow{BC} = -\mathbf{b}$		B1	
aiii		$\overrightarrow{AF} = \frac{2}{3}$ " \overrightarrow{AB} " = $\frac{2}{3}$ ("2 b - a ")		B1 ft	
		(o.e.)			
aiv	$\overrightarrow{FC} = \frac{1}{3}$ "(2 b - a)" +"- b " OR			M1	
	$-\frac{2}{3}"(2\mathbf{b}-\mathbf{a})"-\mathbf{a}+\mathbf{b}$				
		$\overrightarrow{FC} = -\frac{1}{3} (\mathbf{a} + \mathbf{b}) (\text{o.e.})$		A1	
b		$\overrightarrow{FE} = -\frac{\lambda}{3}"(\mathbf{a}+\mathbf{b})"$	1	B1 ft	
с	$\overrightarrow{OE} = \mathbf{a} + \left\ \frac{2}{3}(2\mathbf{b} - \mathbf{a})\right\ + \left\ \left(-\frac{\lambda}{3}(\mathbf{a} + \mathbf{b})\right)\right\ $		2	M1	
	$\left(=\overrightarrow{OA}+\overrightarrow{AF}+\overrightarrow{FE}\right)$				
	OR				
	$\overrightarrow{OE} = \overrightarrow{OC} + \overrightarrow{CE} = \mathbf{b} + (\lambda - 1)'' \left(-\frac{1}{3} (\mathbf{a} + \mathbf{b}) \right)''$				
	$\begin{bmatrix} FE = FC + CE \therefore CE = FE - FC = (\lambda - 1)FC \end{bmatrix}$ (o.e.)				
		$\overrightarrow{OE} = \mathbf{a} \left(\frac{1}{3} - \frac{\lambda}{3} \right) + \mathbf{b} \left(\frac{4}{3} - \frac{\lambda}{3} \right)$ (o.e.)		A1	Accept $\frac{1}{3}\mathbf{a} - \frac{1}{3}\lambda\mathbf{a} + \frac{4}{3}\mathbf{b} - \frac{1}{3}\lambda\mathbf{b}$
d	$\mu''(2\mathbf{b}-\mathbf{a})'' = ''\mathbf{a}\left(\frac{1}{3}-\frac{\lambda}{3}\right) + \mathbf{b}\left(\frac{4}{3}-\frac{\lambda}{3}\right)''$		6	M1	

	Equating components one pair of components			M1	
				(dep)	
		of a : $-\mu = \frac{1}{3} - \frac{\lambda}{3}$		A1	
		of b : $2\mu = \frac{4}{3} - \frac{\lambda}{3}$		A1	
		$\lambda = 2$		A1	
		$\mu = \frac{1}{3}$		A1	
e		Congruent, similar triangles or	1	B1	
		same area			
		$OCE \cong CFB, OCE \square CFB$			
		Stating at least 3 pairs of sides			
		and/or 3 pairs of angles are equal			
					Total 15 marks

11 a	-0.28, 3.28 (awrt)	2	B1	
11 u	0.20, 5.20 (awit)	-	B1	
b	-1 mark for	3	B3	Accuracy for both plotting and
	straight line segments			drawing is $\pm \frac{1}{2}ss$
	each point missed			ft from their table values
	each missed segment			Only penalise straight line
	each point not plotted			segments in the range $-0.5 < x < 1$
	each point incorrectly plotted			and/or $1 < x < 1.5$
	tramlines			
	very poor curve			
с	-0.3 , 1.3	2	B1 B1	For -0.3 accept any value in the range $-0.3 \rightarrow -0.2$
				For 1.3 accept any value in the range $1.2 \rightarrow 1.3$
d	x < -0.7 x > 0.4 x < 1.9	3	B1 ft B1	0.4 < <i>x</i> < 1.9 only stated scores B0, B1, B1
	x < 1.9		ft	Accept weak inequalities
			B1 ft	If none of the B marks are earned, award B1, B0, B0 if and only if all 3 critical values are stated and are
				correct (± small square)

e	$\left(-2x^{3}+3x^{2}+2x\right)+\left(-\frac{1}{2}x-2\right)=0$		6	M1
	$\therefore \left(-2x^3 + 3x^2 + 2x\right) = \frac{1}{2}x + 2 \text{(oe,}$			
	allow 1 sign slip)) 2^{x+2} (out,			
	$y = "\frac{1}{2}x + 2"$ drawn and going			$\begin{array}{c} M1 \\ (dep) \end{array} \text{Accuracy of the line} \pm \frac{1}{2} ss \text{ of the} \end{array}$
	through (0, 2) or (2, 3) extrapolating if necessary) two given points
		Line going through (0, 2) and (2, 3)extrapolating where necessary and intersecting candidate's		A1 2 nd M can imply the first M
		$y = -2x^3 + 3x^2 + 2x$ three times		
		-0.8 , 0.8 ,		A1 ft
		1.6		A1 ft A1
				ft Total 16 marks

Pearson Education Limited. Registered company number 872828 with its registered office at 80 Strand, London, WC2R 0RL, United Kingdom