


1 Electricity is generated in a nuclear power station.

Fission is the process by which energy is released in the nuclear reactor.

(a) **Figure 1** shows the first part of the nuclear fission reaction.

Complete **Figure 1** to show how the fission process starts a chain reaction.


Figure 1



(3)

(b) **Figure 2** shows the inside of a nuclear reactor in a nuclear power station.

Figure 2



In a nuclear reactor a chain reaction occurs, which causes neutrons to be released.

The control rods absorb neutrons.

The control rods can be moved up and down.


Explain how the energy released by the chain reaction is affected by moving the control rods.

.....
.....
.....
.....

(2)

- (c) **Figure 3** shows how the power output of the nuclear reactor would change if the control rods were removed.

Figure 3



Calculate the rate of increase of power output at 10 minutes.

.....

.....

.....

.....

Rate of increase of power output = MW / minute

(2)
(Total 7 marks)

2

Alpha particles, beta particles and gamma rays are types of nuclear radiation.

- (a) Describe the structure of an alpha particle.

.....

.....

(1)

(b) Nuclear radiation can change atoms into ions by the process of ionisation.

(i) Which type of nuclear radiation is the least ionising?

Tick (✓) **one** box.

alpha particles

beta particles

gamma rays

(1)

(ii) What happens to the structure of an atom when the atom is ionised?

.....
.....

(1)

(c) People working with sources of nuclear radiation risk damaging their health.

State **one** precaution these people should take to reduce the risk to their health.

.....
.....

(1)

(d) **In this question you will be assessed on using good English, organising information clearly and using specialist terms where appropriate.**

The type of radiation emitted from a radioactive source can be identified by comparing the properties of the radiation to the properties of alpha, beta and gamma radiation.

Describe the properties of alpha, beta and gamma radiation in terms of their:

- penetration through materials
- range in air
- deflection in a magnetic field.

(6)

(Total 10 marks)

3

(a) Radioactive sources that emit alpha, beta or gamma radiation can be dangerous.

What is a possible risk to health caused by using a radioactive source?


.....
.....

(1)

- (b) In an experiment, a teacher put a 2 mm thick lead sheet in front of a radioactive source. She used a detector and counter to measure the radiation passing through the lead sheet in one minute.

She then put different numbers of lead sheets, each 2 mm thick, in front of the radioactive source and measured the radiation passing through in one minute.

The apparatus the teacher used is shown in **Figure 1**.



- (i) When using a radioactive source in an experiment, how could the teacher reduce the risk to her health?

Suggest **one** way.

.....

.....

(1)

- (ii) The number recorded on the counter is actually higher than the amount of radiation detected from the source.


Complete the following word equation.

| | | | | |
|------------------------------------|---|--|---|-----------------|
| The number recorded on the counter | = | The amount of radiation detected from the source | + | radiation |
|------------------------------------|---|--|---|-----------------|

(1)

- (c) The readings taken by the teacher are plotted in **Figure 2**.

Figure 2



- (i) Draw a line of best fit to complete **Figure 2**. (1)
- (ii) How does the amount of radiation **absorbed** by the lead change as the total thickness of the lead is increased?

 (1)
- (iii) Use **Figure 2** to estimate the reading on the counter when the total thickness of the lead is increased to 12 mm.
 Estimated counter reading = (1)

(d) What type of radiation was emitted from the radioactive source?

Draw a ring around the correct answer.

alpha

beta

gamma

Give a reason for your answer.

.....
.....

(2)
(Total 8 marks)

4

Different radioactive isotopes have different values of half-life.


(a) What is meant by the 'half-life' of a radioactive isotope?

.....
.....
.....

(1)

(b) **Figure 1** shows how the count rate from a sample of a radioactive isotope varies with time.

Figure 1



Use information from **Figure 1** to calculate the half-life of the radioactive isotope.

Show clearly on **Figure 1** how you obtain your answer.

Half-life = days

(2)

(c) The table below shows data for some radioactive isotopes that are used in schools.

| Radioactive isotope | Type of radiation emitted | Half-life in years |
|---------------------|---------------------------|----------------------|
| Americium-241 | Alpha and gamma | 460 |
| Cobalt-60 | Gamma | 5 |
| Radium-226 | Alpha, beta and gamma | 1600 |
| Strontium-90 | Beta | 28 |
| Thorium-232 | Alpha and beta | 1.4×10^{10} |

(i) State which radioactive isotope in the table above emits only radiation that is **not** deflected by a magnetic field.

Give a reason for your choice.

.....

.....


.....

.....

(2)

- (ii) **Figure 2** shows a radioactive isotope being used to monitor the thickness of paper during production.

Figure 2



State which radioactive isotope in the table should be used to monitor the thickness of the paper.

Explain your choice.

.....

.....

.....

.....

.....

(3)

All the radioactive isotopes in the table have practical uses.

State which source in the table would need replacing most often.

Explain your choice.

.....

.....

.....

.....

.....


(3)

- (iii) When the radioactive isotopes are not in use, they are stored in lead-lined wooden boxes.

The boxes reduce the level of radiation that reaches the surroundings.

Figure 3 shows two of these boxes.

Figure 3



© David McKean

State **one** source from the table which emits radiation that could penetrate the box.

Explain your answer.

.....

.....

.....

.....

.....

.....

(3)
(Total 14 marks)

5

Nuclear fission and nuclear fusion are two processes that release energy.

(a) (i) Use the correct answer from the box to complete each sentence.

| | | |
|----------------|-----------------|------|
| Geiger counter | nuclear reactor | star |
|----------------|-----------------|------|

Nuclear fission takes place within a

Nuclear fusion takes place within a

(2)

(ii) State **one** way in which the process of nuclear fusion differs from the process of nuclear fission.

.....
.....

(1)

(b) The following nuclear equation represents the fission of uranium-235 (U-235).



Chemical symbols:

Ba - barium

Kr - krypton


(i) Use the information in the equation to describe the process of nuclear fission.

.....
.....
.....
.....
.....
.....
.....
.....
.....

(4)

- (ii) An isotope of barium is Ba-139.
Ba-139 decays by beta decay to lanthanum-139 (La-139).


Complete the nuclear equation that represents the decay of Ba-139 to La-139.



(3)
(Total 10 marks)

6

A teacher used the equipment shown in the diagram to measure the count rate at different distances from a radioactive source.



- (a) Her results are shown in **Table 1**.

Table 1

| Distance in metres | Count rate in counts per minute | Corrected count rate in counts per minute |
|--------------------|---------------------------------|---|
| 0.4 | 143 | 125 |
| 0.6 | 74 | 56 |
| 0.8 | 49 | 31 |
| 1.0 | 38 | 20 |
| 1.2 | 32 | 14 |
| 1.4 | 28 | 10 |
| 1.6 | 18 | 0 |
| 1.8 | 18 | 0 |
| 2.0 | 18 | 0 |

The background count rate has been used to calculate the corrected count rate.

- (i) What is the value of the background count rate?

Background count rate = counts per minute

(1)

(ii) What information does the corrected count rate give?

.....
.....

(1)

(iii) The radioactive source used in the demonstration emits only one type of radiation.

The radioactive source is **not** an alpha emitter.


How can you tell from the data in the table?

.....
.....

(1)

(iv) Plot a graph of corrected count rate against distance for distances between 0.4 m and 1.4 m.

Draw a line of best fit to complete the graph.



(3)

- (v) The 'half-distance' is the distance a detector has to be moved away from a radioactive source for the corrected count rate to halve.

A student has the hypothesis:

A radioactive source has a constant 'half-distance'.

Table 1 has been repeated for your information.

Table 1

| Distance in metres | Count rate in counts per minute | Corrected count rate in counts per minute |
|--------------------|---------------------------------|---|
| 0.4 | 143 | 125 |
| 0.6 | 74 | 56 |
| 0.8 | 49 | 31 |
| 1.0 | 38 | 20 |
| 1.2 | 32 | 14 |
| 1.4 | 28 | 10 |
| 1.6 | 18 | 0 |
| 1.8 | 18 | 0 |
| 2.0 | 18 | 0 |

Use **Table 1** to determine if the hypothesis is correct for this radioactive source.

You should use calculations in your answer.

.....

.....

.....

.....

.....

.....


.....

.....


(3)

(b) A teacher places a beta source and a detector in a magnetic field.

The arrangement of the magnetic field is shown.



.....The teacher repeated the experiment with the magnetic field in a different direction.



A set of results is shown in **Table 2**.

Table 2

| Distance between source and detector in metres | Count rate in counts per minute without magnetic field | Count rate in counts per minute in Experiment 1 | Count rate in counts per minute in Experiment 2 |
|--|--|---|---|
| 0.8 | 48 | 48 | 32 |

(i) Describe **and** explain the effect of the magnetic field on the count rate detected by the detector.

.....

.....

.....

.....

.....

.....

.....

(2)

- (ii) The experiment is repeated with a different distance between the source and the detector.

Table 3 shows the repeated results.

Table 3

| Distance between source and detector in metres | Count rate in counts per minute without magnetic field | Count rate in counts per minute in Experiment 1 | Count rate in counts per minute in Experiment 2 |
|--|--|---|---|
| 1.8 | 19 | 18 | 20 |

Explain these results.

.....

.....

.....

.....

.....

.....

(2)
(Total 13 marks)

7

- (a) Sources of background radiation are either natural or man-made.

Which **two** of the sources listed in the box are *natural* sources of background radiation?


Draw a ring around each of your answers.

| | | | |
|-------------|-------------------|--------|-----------|
| cosmic rays | nuclear accidents | X-rays | radon gas |
|-------------|-------------------|--------|-----------|

(2)

- (b) A teacher used a Geiger-Müller (GM) tube and counter to measure the background radiation in her laboratory. The teacher reset the counter to zero, waited one minute and then took the count reading. The teacher repeated this two more times.

The three readings taken by the teacher are given in the table.



- (i) The three readings are different.

What is the most likely reason for this?

Tick (✓) **one** box.

The teacher did not reset the counter to zero.

Radioactive decay is a random process.

The temperature in the laboratory changed.

(1)

- (ii) Calculate the mean (average) value of the three readings given in the table.


.....

Mean (average) value = counts

(1)

- (iii) The diagram shows how the teacher used the GM tube and counter to measure the radiation emitted from a radioactive source.

The counter was reset to zero. The count after one minute was 159.



Calculate how many counts were due to the radiation from the radioactive source.


.....

Counts due to the radiation from the radioactive source =

(1)

- (iv) The teacher then put a powerful magnet between the radioactive source and the GM tube.

The counter was reset to zero. The number on the counter shows the count after one minute.



What type of radiation was being emitted from the radioactive source?

Draw a ring around your answer.

alpha

beta


gamma

Explain the reason for your answer.

.....

(3)

- (c) At the end of the lesson the teacher put the radioactive source back inside its storage box.



Why is the inside of the box lined with lead?

.....
.....

(1)

- (d) Which **one** of the following questions **cannot** be answered by scientific study?

Tick (✓) **one** box.

Where does background radiation come from?

What is meant by the half-life of a radioactive source?


Should radioactive waste be dumped in the oceans?

(1)
(Total 10 marks)

8

(a) A teacher used a Geiger-Müller (GM) tube and counter to measure the *background radiation* in her laboratory.

The teacher reset the counter to zero, waited one minute and then took the count reading. The teacher repeated the procedure two more times.



(i) Background radiation can be either from natural sources or from man-made sources.

Name **one man-made** source of background radiation.

.....

(1)

(ii) The three readings taken by the teacher are given in the table.

| Count after one minute |
|------------------------|
| 15 |
| 24 |
| 18 |

The readings given in the table are correct.

Why are the readings different?

.....

.....

(1)

- (b) Some scientists say they have found evidence to show that people living in areas of high natural background radiation are less likely to develop cancer than people living in similar areas with lower background radiation.

The evidence these scientists found does not definitely mean that the level of background radiation determines whether a person will develop cancer.

Suggest a reason why.

.....
.....

(1)

- (c) An atom of the isotope radon-222 emits an alpha particle and decays into an atom of polonium.

An alpha particle is the same as a helium nucleus. The symbol below represents an alpha particle.



- (i) How many protons and how many neutrons are there in an alpha particle?

Number of protons =

Number of neutrons =

(2)


- (ii) The decay of radon-222 can be represented by the equation below.

Complete the equation by writing the correct number in each of the **two** boxes.



(2)

- (d) The graph shows how, in a sample of air, the number of radon-222 nuclei changes with time.



Use the graph to find the half-life of radon-222.

Show clearly on the graph how you obtain your answer.

Half-life = days

(2)
(Total 9 marks)

9

A doctor uses the radioactive isotope technetium-99 to find out if a patient's kidneys are working correctly.




The doctor injects a small amount of technetium-99 into the patient's bloodstream. Technetium-99 emits gamma radiation.

If the patient's kidneys are working correctly, the technetium-99 will pass from the bloodstream into the kidneys and then into the patient's urine.

Detectors are used to measure the radiation emitted from the kidneys.

The level of radiation emitted from each kidney is recorded on a graph.



(a) How do the graphs show that technetium-99 is passing from the bloodstream into each kidney?

.....
.....

(1)

- (b) By looking at the graphs, the doctor is able to tell if there is a problem with the patient's kidneys.

Which **one** of the following statements is correct?

Put a tick (✓) in the box next to your answer.

Only the right kidney is working correctly.

Only the left kidney is working correctly.

Both kidneys are working correctly.

Explain the reason for your answer.

.....

.....


.....

.....

(3)
(Total 4 marks)

10

The pie chart shows the average proportions of background radiation from various sources in the UK.



- (a) Three sources of background radiation are given in **List A**. Statements about sources of background radiation are given in **List B**.

Draw **one** line to link each source of background radiation in **List A** to the statement about that source given in **List B**.

Draw only **three** lines.

List A

X-rays

Cosmic rays

Radon gas

List B

Are used to show broken bones.

The radiation comes from outer space.


Comes from soil containing a radioactive isotope of potassium.

On average gives 50% of all background radiation.

(3)

- (b) The level of background radiation from cosmic rays is not the same everywhere. For every 30 metres above sea level, the amount of background radiation increases by one unit.

The diagram shows the position of two villages, **A** and **B**, built on a hill.



How is the amount of background radiation from cosmic rays different in village **A** compared to village **B**?

To obtain full marks, you must include a calculation in your answer.

.....

.....

.....

.....


.....

.....

(3)
(Total 6 marks)

11

The pie chart shows the sources of the background radiation and the radiation doses that the average person in the UK is exposed to in one year. Radiation dose is measured in millisieverts (mSv).



(a) (i) What is the total radiation dose that the average person in the UK receives?

.....
.....

Total radiation dose = mSv

(1)

(ii) A student looked at the pie chart and then wrote down three statements.

Which **one** of the following statements is a correct conclusion from this data?

Put a tick (✓) in the box next to your answer.

In the future, more people will be exposed to a greater proportion of radon gas.

People that have never had an X-ray get 50 % of their radiation dose from radon gas.

The radiation dose from natural sources is much greater than from artificial sources.

(1)

- (b) The concentration of radon gas inside a home can vary from day to day.

The table gives data for the radiation measured in homes in four different parts of the UK. The radiation was measured using two detectors, one in the living room and one in the bedroom. The measurements were taken over 3 months.

| Area of the UK | Number of homes in the area | Number of homes in the sample | Average radiation in Bq/m ³ | Maximum radiation in Bq/m ³ |
|----------------|-----------------------------|-------------------------------|--|--|
| A | 590 000 | 160 | 15 | 81 |
| B | 484 000 | 130 | 18 | 92 |
| C | 221 000 | 68 000 | 162 | 10 000 |
| D | 318 000 | 35 300 | 95 | 6 900 |

- (i) Give **one** reason why the measurements were taken over 3 months using detectors in different rooms.

.....

(1)

- (ii) Use information from the table to suggest why a much higher proportion of homes were sampled in areas **C** and **D** than in areas **A** and **B**.

.....

(2)
 (Total 5 marks)

12

The table shows the average background radiation dose from various sources that a person living in the UK receives in one year.

| Source of background radiation | Average radiation dose received each year in dose units |
|---------------------------------------|---|
| Cosmic rays (from space) | 300 |
| Food and drink | 250 |
| Medical treatments (including X-rays) | 350 |
| Radon gas | 1250 |
| Rocks | 350 |
| TOTAL | 2500 |

(a) (i) A student looked at the data in the table and then wrote down four statements.

Only **two** of the statements are true.

Put a tick (✓) in the boxes next to the **two** true statements.

More than half of the average radiation dose comes from radon gas.

On average, cosmic rays produce less background radiation than rocks.

Everyone living in the UK receives the same background radiation dose.

Having no X-rays reduces a person's radiation dose.

(2)

(ii) Each time a chest X-ray is taken, the patient receives about 100 units of radiation.

How many chest X-rays would just exceed the yearly average dose for medical treatments?


.....

Number of chest X-rays =

(2)

(b) Exposure to radiation can cause cancer.

The graphs, **A**, **B** and **C**, show three different ways that the exposure to radiation and the risk of getting cancer could be linked.



(i) What do all three of these graphs suggest happens to the risk of getting cancer when the radiation dose goes from moderate to high?

.....

.....

(1)

- (ii) Some scientists believe that exposure to **low** radiation doses reduces the chance that a person will get cancer. This effect is called 'radiation hormesis'.

Which one of the graphs, **A**, **B** or **C**, shows 'radiation hormesis'?

Write your answer in the box.

Give a reason for your answer.

.....

.....

.....

(2)

- (c) Scientists did an experiment in which mice were exposed to different doses of radiation.

The results from the experiment are given in the table.

| Description of exposure | Percentage of mice getting cancer |
|--|-----------------------------------|
| Mice exposed to a low dose of radiation and then a high dose of radiation. | 16% |
| Mice exposed to a high dose of radiation only. | 46% |

- (i) Do the results from this experiment provide evidence to support 'radiation hormesis'?

Draw a ring around your answer.

NO

YES

Explain the reason for your answer.

.....

.....

.....

.....

.....

(2)

- (ii) Complete the following sentence by drawing a ring around the correct word in the box.

Using animals in scientific experiments raises

| |
|---------------|
| environmental |
| ethical |
| social |

issues.


(1)
(Total 10 marks)

13

The pie chart shows the sources of the background radiation and the radiation doses that the average person in the UK is exposed to in one year.

Radiation dose is measured in millisieverts (mSv).

Other sources, including nuclear weapons testing,
nuclear accidents and power stations
0.01 mSv



- (a) (i) What is the radiation dose that the average person in the UK receives from radon gas?

.....
.....

Radiation dose from radon gas = mSv

(1)

- (ii) A person may receive a higher than average dose of radiation from background sources.

Suggest **two** reasons why.

1

.....

2

.....

(2)

- (b) Exposure to radon gas can cause lung cancer.
A recent study has compared the risk of getting lung cancer, by the age of 75 years, for cigarette smokers and non-smokers.
The people in the study had been exposed throughout their lives to different levels of radon gas.
A summary of the data produced from the study is given in the table.

| Exposure to radon gas | Risk of lung cancer by age of 75 | |
|-----------------------|----------------------------------|--------|
| | Non-smoker | Smoker |
| No exposure | 0.4 % | 10 % |
| Moderate exposure | 1.0 % | 14 % |
| Very high exposure | 1.5 % | 32 % |

- (i) Why were people that have had **no exposure** to radon gas included in the study?

.....

.....

(1)

- (ii) Using information from the table, what conclusions can be made about exposure to radon gas and the risk of getting lung cancer?

.....

.....

.....


.....

.....

(2)

- (c) At the moment, the regulations designed to protect people from over-exposure to radiation are based on a model called the 'linear no-threshold' (LNT) model. Some scientists believe that the LNT model is too simple. These scientists believe that at low radiation levels a process called 'radiation hormesis' happens.

The graphs show that each model suggests a link between the risk of developing a cancer and exposure to low levels of radiation.



The link between the risk of developing cancer and exposure to low levels of radiation suggested by each of the models is different.

Describe how.

.....

.....

.....

.....

.....

(2)

- (d) Scientists have conducted experiments in which mice have been exposed to different levels of radiation. The number of mice developing a cancer has then been measured.

Discuss whether it is ethical to use animals in scientific experiments.

.....

.....

.....

.....

.....

(2)
(Total 10 marks)

14

Food irradiation is a process that exposes food to radiation. Irradiation can be used to kill the bacteria that cause food poisoning or to slow down the ripening of fresh fruit and vegetables. Frozen foods and food inside packaging can also be irradiated.

(a) The table gives information about five radioactive isotopes.

| Isotope | Half-life | Radiation emitted |
|---------------|-----------|-------------------|
| Caesium-134 | 2.1 years | beta |
| Cobalt-60 | 5.3 years | gamma |
| Curium-242 | 160 days | alpha |
| Strontium-90 | 28 years | beta |
| Technetium-99 | 6 hours | gamma |

Which of these radioactive isotopes would be most suitable for irradiating food?

.....

Explain the reasons for your choice.

.....

.....

.....

.....

.....

(3)

(b) Many people think that food should not be irradiated. Consumer groups have said that they are worried about the nutritional value and safety of eating irradiated foods.

(i) Suggest **one** reason why some people may be concerned about the safety of eating irradiated food.

.....

.....

(1)

- (ii) Independent scientific committees in several countries, including Sweden, Canada and the UK, have concluded that it is safe to eat irradiated food.

These scientific committees need to be independent from government influence.

Suggest why.

.....
.....

(1)

- (iii) One group of scientists has compared the vitamin content of non-irradiated foods with irradiated foods.

The table below gives the data obtained for 1 kg of cooked chicken.

| Vitamin | Non-irradiated food in milligrams | Irradiated food in milligrams |
|----------------|--|--|
| B6 | 1.22 | 1.35 |
| B12 | 21.00 | 28.00 |
| E | 3.30 | 2.15 |
| Niacin | 58.00 | 55.50 |
| Riboflavin | 2.10 | 2.25 |

Considering only the data in the table, is it valid to conclude that irradiated food is less nutritional than non-irradiated food?

Explain your answer.

.....
.....
.....
.....
.....

(2)

- (iv) In a restaurant, meals with ingredients that have been irradiated must be clearly identified on the menu.

It is important that people eating in a restaurant are given this information.

Suggest why.

.....
.....

(1)

- (c) The isotope caesium-137 decays by emitting beta radiation.
Caesium-137 has a half-life of 30 years.

- (i) What is a beta particle, and from which part of an atom is a beta particle emitted?

.....
.....

(1)

- (ii) A sample containing caesium-137 has a count rate of 600 counts per minute.

Calculate how long it would take for the count rate from the sample to fall to 75 counts per minute.

Show clearly how you work out your answer.

.....
.....
.....


Time taken = years

(2)

(Total 11 marks)

15

The pie chart shows the average proportions of natural background radiation from various sources in the UK.



(a) (i) Complete the following sentence.

On average, of the natural background radiation in the UK comes from radon gas.

(1)

- (ii) Radon gas is found inside homes.

The table shows the results from measuring the level of radon gas inside four homes in one area of the UK.

| Home | Level of radon gas in Bq per m ³ of air |
|------|--|
| 1 | 25 |
| 2 | 75 |
| 3 | 210 |
| 4 | 46 |
| Mean | 89 |

One of the homes has a much higher level of radon gas than the other three homes.

What should be done to give a more reliable mean for the homes in this area of the UK?

Put a tick (✓) in the box next to your answer.

ignore the data for home number 3

measure the radon gas level in more homes in this area

include data for homes from different areas of the UK

(1)

- (b) Each atom of radon has 86 protons and 136 neutrons.

- (i) How many electrons does each atom of radon have?

Draw a ring around your answer.

50

86

136

222

(1)

(ii) How many particles are there in the nucleus of a radon atom?

Draw a ring around your answer.

50

86

136

222

(1)
(Total 4 marks)

16

(a) Background radiation is all around us all the time.

(i) Radon is a natural source of background radiation.

Name another natural source of background radiation.

.....

(1)

(ii) X-rays are an artificial source of background radiation.


Name another artificial source of background radiation.

.....

(1)

(iii) An atom of radon-222 decays by emitting an alpha particle.

The equation representing the decay is shown below.



How can you tell from the equation that 'X' is not an atom of radon?

.....

.....

(1)

(b) Having an X-ray taken increases your exposure to radiation.

The table gives:

- the radiation doses received for 6 different medical X-rays;
- the number of days' of exposure to natural background radiation each dose is equivalent to.

| Medical X-ray | Radiation dose received (in arbitrary units) | Equivalent number of days of exposure to natural background radiation |
|----------------------|---|--|
| Chest | 2 | 2.4 |
| Skull | 7 | 8.4 |
| Pelvis | 22 | 26.4 |
| Hip | 44 | 52.8 |
| Spine | 140 | |
| CT head scan | 200 | 240 |

A hospital patient has an X-ray of the spine taken.

Calculate the number of days of exposure to natural background radiation that an X-ray of the spine is equivalent to.

Show how you work out your answer.







.....
.....
.....

Equivalent number of days =

(2)

- (c) Scientists have shown that X-rays increase the risk of developing cancer. The scientists came to this conclusion by studying the medical history of people placed in one of two groups, **A** or **B**. The group into which people were put depended on their X-ray record.

- (i) Person **J** has been placed into group **A**. Place each of the people, **K**, **L**, **M**, **N** and **O**, into the appropriate group, **A** or **B**.

| | | | | | | |
|-----------------------------|---|---|---|---|---|---|
| Person | J  | K  | L  | M  | N  | O  |
| Medical X-ray record | 3 arm | None | None | 2 skull | None | 4 leg |

| Group A | Group B |
|----------------|----------------|
| J | |

(1)

- (ii) To be able to make a fair comparison, what is important about the number of people in each of the two groups studied by the scientists?

.....

(1)

- (iii) What data would the scientists have compared in order to come to the conclusion that X-rays increase the risk of developing cancer?

.....

(1)

- (iv) The chance of developing cancer due to a CT head scan is about 1 in 10 000.
The chance of developing cancer naturally is about 1 in 4.

A hospital patient is advised by a doctor that she needs to have a CT head scan.
The doctor explains to the patient the risks involved.

Do you think that the patient should give her permission for the CT scan to be taken?

Draw a ring around your answer.

Yes

No

Give a reason for your answer.


.....

.....

(1)
(Total 9 marks)

17

The pie chart shows the average proportions of background radiation from various sources in the UK.



Three sources of background radiation are given in **List A**.
Statements about sources of background radiation are given in **List B**.

Draw **one** line to link each source of background radiation in **List A** to the statement about that source given in **List B**.

Draw only **three** lines.

List A

List B

X-rays

Cosmic rays

Radon gas

Are used to show broken bones.

The radiation comes from outer space.


Comes from soil containing a radioactive isotope of potassium.

Gives about 50 % of all background radiation.

(Total 3 marks)

18

(a) A doctor uses the radioactive isotope technetium-99 to find out if a patient's kidneys are working correctly.



The doctor injects a small amount of technetium-99 into the patient's bloodstream.

Technetium-99 emits *gamma radiation*.

Give **two** reasons why an isotope that emits gamma radiation is injected into the patient rather than an isotope that emits alpha radiation.


- 1.....
-
- 2.....
-

(2)

- (b) If the patient's kidneys are working correctly, the technetium-99 will pass from the bloodstream into the kidneys and then into the patient's urine.

Detectors are used to measure the radiation emitted from the kidneys.

The level of radiation emitted from each kidney is recorded on a graph.



- (i) How do the graphs show that technetium-99 is passing from the bloodstream into each kidney?

.....

(1)

- (ii) By looking at the graphs, the doctor is able to tell if there is a problem with the patient's kidneys.

Which **one** of the following statements is correct?

Put a tick (✓) in the box next to your answer.

Only the right kidney is working correctly.

Only the left kidney is working correctly.

Both kidneys are working correctly.

Explain the reason for your answer.

.....

(3)

(c) The patient was worried about having a radioactive isotope injected into their body. The doctor explained that the risk to the patient's health was very small as technetium-99 has a short *half-life*.

(i) What does the term *half-life* mean?

.....
.....

(1)

(ii) Explain why it is important that the doctor uses an isotope with a short half-life rather than an isotope with a long half-life.


.....
.....
.....
.....

(2)

(Total 9 marks)

19

The pie chart shows the sources of the background radiation and the radiation doses that the average person in the UK is exposed to in one year. Radiation dose is measured in millisieverts (mSv).



(a) (i) What is the total radiation dose that the average person in the UK receives?

.....
.....

Total radiation dose = mSv

(1)

(ii) A student looked at the pie chart and then wrote down three statements.

Which **one** of the following statements is a correct conclusion from this data?

Put a tick (✓) in the box next to your answer.

In the future, more people will be exposed to a greater proportion of

radon gas.

People that have never had an X-ray get 50% of their radiation dose from

radon gas.

The radiation dose from natural sources is much greater than from artificial

sources.

(1)

(b) The concentration of radon gas inside a home can vary from day to day. In some homes, the level can build up to produce a significant health risk. It is estimated that each year 1000 to 2000 people die because of the effects of radiation from radon gas.

(i) It is not possible to give an exact figure for the number of deaths caused by the effects of radiation from radon gas. Why?

.....
.....

(1)

The table gives data for the radiation levels measured in homes in 4 different parts of the UK. The radiation levels were measured using two detectors, one in the living room and one in the bedroom. The measurements were taken over 3 months.

| Area of the UK | Number of homes in the area | Number of homes in the sample | Average radiation level in Bq/m ³ | Maximum radiation level in Bq/m ³ |
|----------------|-----------------------------|-------------------------------|--|--|
| A | 590 000 | 160 | 15 | 81 |
| B | 484 000 | 130 | 18 | 92 |
| C | 221 000 | 68 000 | 162 | 10 000 |
| D | 318 000 | 35 300 | 95 | 6 900 |

- (ii) Give **one** reason why the measurements were taken over 3 months using detectors in different rooms.

.....

(1)

- (iii) Use information from the table to suggest why a much higher proportion of homes were sampled in areas **C** and **D** than in areas **A** and **B**.


.....

(2)

(Total 6 marks)

20

- (a) The pie chart shows the average proportions of natural background radiation from various sources in one part of the UK.



- (i) What proportion of the background radiation comes from radon gas?

.....

(1)

- (ii) Suggest why our bodies are slightly radioactive.


.....

.....

(1)

- (b) The level of background radiation from cosmic rays is not the same everywhere. For every 30 metres above sea level, the amount of background radiation increases by one unit.

The diagram shows the position of two villages, **A** and **B**, built on a hill.



How is the amount of background radiation from cosmic rays different in village **A** compared to village **B**?

To obtain full marks you must include a calculation in your answer.

.....

.....

.....

.....


.....

.....

(3)
(Total 5 marks)

21

A radioactive source emits alpha (α), beta (β) and gamma (γ) radiation. The diagram shows what happens to the radiation as it passes between two charged metal plates.



(a) Which line **P**, **Q** or **R** shows the path taken by:


(i) alpha radiation

(1)

(ii) gamma radiation?

(1)


- (b) The diagram shows three different boxes and three radioactive sources. Each source emits only one type of radiation and is stored in a different box. The box reduces the amount of radiation getting into the air.



Draw **three** lines to show which source should be stored in which box so that the minimum amount of radiation gets into the air.

(2)

- (c) The graphs show how the count rates from three different radioactive sources, **J**, **K**, and **L**, change with time.



- (i) Which source, **J**, **K**, or **L**, has the highest count rate after 24 hours?
 (1)
- (ii) For source **L**, what is the count rate after 5 hours?
 counts per second (1)
- (iii) Which source, **J**, **K**, or **L**, has the longest half-life?
 (1)
- (iv) A radioactive source has a half-life of 6 hours.
 What might this source be used for?
 Put a tick (✓) in the box next to your choice.
- To monitor the thickness of paper as it is made in a factory
- To inject into a person as a medical tracer
- To make a smoke alarm work

(1)
 (Total 8 marks)

22

- (a) A radioactive source emits alpha (α), beta (β) and gamma (γ) radiation.
- (i) Which **two** types of radiation will pass through a sheet of card?
 (1)
- (ii) Which **two** types of radiation would be deflected by an electric field?
 (1)
- (iii) Which type of radiation has the greatest range in air?
 (1)

- (b) A student suggests that the radioactive source should be stored in a freezer at $-20\text{ }^{\circ}\text{C}$. The student thinks that this would reduce the radiation emitted from the source.

Suggest why the student is wrong.

.....

(1)


- (c) Phosphorus-32 is a radioactive isotope that emits beta radiation.

- (i) How is an atom of phosphorus-32 different from an atom of the stable isotope phosphorus-31?

.....

(1)

- (ii) The graph shows how the count rate of a sample of phosphorus-32 changes with time.



Use the graph to calculate the half-life of phosphorus-32.


Show clearly how you used the graph to obtain your answer.

.....

Half-life = days

(2)

- (iii) Plants use phosphorus compounds to grow. Watering the root system of a plant with a solution containing a phosphorus-32 compound can help scientists to understand the growth process.



Explain why phosphorus-32 is suitable for use as a tracer in this situation.

.....

.....

.....

.....

(2)
(Total 9 marks)

23

The table shows the average background radiation dose from various sources that a person living in Britain receives in one year.

| Source of background radiation | Average amount each year in dose units |
|---------------------------------------|--|
| Buildings | 50 |
| Food and drink | 300 |
| Medical treatments (including X-rays) | 300 |
| Radon gas | 1250 |
| Rocks | 360 |
| Space (cosmic rays) | 240 |
| TOTAL | 2500 |

(a) Only **two** of the following statements are true.

Tick (✓) the boxes next to the true statements.

Half the average background radiation dose comes from radon gas.

Everyone receives the same background radiation dose.

Cosmic rays produce less background radiation than food and drink.

(1)

(b) Most sources of background radiation are natural but some are artificial (man-made).

Which source of background radiation given in the table is artificial?

.....

(1)

(c) Each time a dental X-ray is taken, the patient receives about 20 units of radiation.

How many dental X-rays would give the yearly average dose for medical treatments?

.....

.....

Number of X-rays =

(2)

(Total 4 marks)

24

- (a) The names of three types of nuclear radiation are given in **List A**. Some properties of these three types of radiation are given in **List B**.

Draw a straight line to link each type of radiation in **List A** to its correct property in **List B**.
Draw only three lines.

| List A | List B |
|---|--|
| Type of nuclear radiation | Property of radiation |
| <div style="border: 1px solid black; width: 60px; height: 40px; margin: 0 auto; display: flex; align-items: center; justify-content: center;">alpha</div> | <div style="border: 1px solid black; width: 280px; height: 40px; margin: 0 auto; display: flex; align-items: center; justify-content: center;">not deflected by an electric field</div> |
| <div style="border: 1px solid black; width: 60px; height: 40px; margin: 0 auto; display: flex; align-items: center; justify-content: center;">beta</div> | <div style="border: 1px solid black; width: 280px; height: 40px; margin: 0 auto; display: flex; align-items: center; justify-content: center;">stopped by thin metal but not paper</div> |
| <div style="border: 1px solid black; width: 60px; height: 40px; margin: 0 auto; display: flex; align-items: center; justify-content: center;">gamma</div> | <div style="border: 1px solid black; width: 280px; height: 40px; margin: 0 auto; display: flex; align-items: center; justify-content: center;">the most strongly ionising</div> |
| | <div style="border: 1px solid black; width: 280px; height: 40px; margin: 0 auto; display: flex; align-items: center; justify-content: center;">will not harm living cells</div> |

(3)

- (b) Nuclear radiation is given out from the centre of some types of atom.

What name is given to the centre of an atom?

(1)

- (c) One of the substances in the table is used as a radioactive tracer. A hospital patient breathes in air containing the tracer. The radiation given out is measured by a doctor using a detector outside the patient's body.

| Substance | Radiation given out | Solid, liquid or gas |
|-----------|---------------------|----------------------|
| X | alpha | gas |
| Y | gamma | gas |
| Z | gamma | solid |

Which **one** of the substances, **X**, **Y** or **Z**, should be used as the tracer?

Give **two** reasons for your answer.

1

.....

2

.....

(3)

- (d) Radiation can also be used to kill the bacteria on fresh food.

Give **one** reason why farmers, shop owners or consumers may want food to be treated with radiation.

.....

.....

(1)

(Total 8 marks)

25

In 1986, a nuclear reactor exploded in a power station at Chernobyl in the Ukraine.

- (a) The table gives information about some of the radioactive substances released into the air by the explosion.

| Radioactive substance | Half-life | Type of radiation emitted |
|-----------------------|-----------|---------------------------|
| Iodine-131 | 8 days | beta and gamma |
| Caesium-134 | 2 years | beta |
| Caesium-137 | 30 years | beta |

- (i) How is the structure of a caesium-134 atom different from the structure of a caesium-137 atom?

.....

(1)

- (ii) What is a beta particle and from which part of an atom is a beta particle emitted?

.....

.....

(1)

- (iii) Once a radioactive substance is dissolved in rainwater, it can enter the food chain.

Following the Chernobyl explosion, some milk supplies were found to be radioactive.

If one litre of milk contaminated with iodine-131 gives a count rate of 400 counts/second, how long will it take for the count rate to fall to 25 counts/second?

Show clearly how you work out your answer.

.....

.....

.....

Time taken = days

(2)

- (iv) After 20 years, the caesium-137 emitted into the atmosphere is a more serious problem than the iodine-131.

Explain why.

.....

.....


.....

.....

.....

(2)

- (b) The bar chart compares the incidence of thyroid cancer in Ukrainian children, aged 0–14 years, before and after the Chernobyl explosion.



Of the children that developed thyroid cancer, 64% lived in the areas most contaminated by the radiation.

Considering this data, can you be certain that a child who developed thyroid cancer between 1986 and 1990 did so because of the Chernobyl explosion?

Explain the reason for your answer.

.....
.....
.....
.....

(2)

- (c) In 1991, some scientists compared the health of two groups of people: a *control* group and a group that had been exposed to the radiation from Chernobyl.

What people would have been in the *control* group?

.....

(1)

- (d) Although there are some risks associated with nuclear power stations, it is likely that new ones will be built.

Give **two** reasons to justify the use of nuclear power.

1

.....

2

.....

(2)

(Total 11 marks)

26

(a) The table gives information about the radioactive isotope, radon-222.

| | |
|-------------------|----------------|
| mass number | 222 |
| atomic number | 86 |
| radiation emitted | alpha particle |

(i) Complete the following sentence.

The mass number is the total number of and
..... inside an atom.

(2)

(ii) Radon-222 is an isotope of radon.

How many protons are there in an atom of radon-222?

.....

(1)

(iii) When an atom of radon-222 emits an alpha particle, the radon-222 changes into an atom of polonium-218.

An alpha particle consists of 2 protons and 2 neutrons.


How is the structure of the nucleus of a polonium-218 atom different from the structure of the nucleus of a radon-222 atom?

.....

(1)

- (b) The pie chart shows the average radiation dose that a person in the UK receives each year from natural background radiation.

The doses are measured in millisieverts (mSv).



- (i) Calculate the proportion of natural background radiation that comes from radon. Show clearly how you work out your answer.

.....
.....

Proportion of radon =

(2)


- (ii) Not all background radiation is from natural sources.

Name **one** source of background radiation that is not natural.

.....

(1)

- (c) The bar chart shows the average yearly dose from natural background radiation in different European countries.



- (i) How many times bigger is the average annual background dose in Germany compared to the UK?

.....

(1)

(ii) The following table gives the effects of different radiation doses on the human body.

| Radiation dose in mSv | Effects |
|-----------------------|---|
| 10 000 | Immediate illness; death within a few weeks |
| 1 000 | Radiation sickness; unlikely to cause death |
| 50 | Lowest dose with evidence of causing cancer |

A family goes to Germany for a two-week holiday. Should they be concerned about the higher level of background radiation in Germany?

Draw a ring around your answer.

Yes **No**

Explain your answer.

.....

.....

.....

.....

(2)
(Total 10 marks)

27

- (a) The names of three types of radiation are given in **List A**. Various properties of these three types of radiation are given in **List B**.

Draw a line to link each type of radiation in **List A** to its correct property in **List B**. Draw only **three** lines.

| List A Type of radiation | List B Property of radiation |
|-----------------------------|---------------------------------|
| alpha (α) | not dangerous |
| beta (β) | stopped by paper |
| gamma (γ) | travels at 300 000 000 m/s |
| | travels up to 1 metre in air |

(3)

- (b) This sign warns people that a radioactive source is being used in a laboratory.



Why is it important to warn people that a radioactive source is being used?

.....
.....

(1)


- (c) To study the blood flow in a patient's lungs, a doctor injects some technetium-99 compound into the patient. The gamma radiation given out by the technetium-99 atoms is detected using a gamma camera outside the patient's body.

Which statement gives the reason why gamma radiation is used? Put a tick (\checkmark) in the box next to your choice.

- It can travel through a vacuum.
- It is not affected by a magnet.
- It can pass through the human body.

(1)

- (d) The graph shows how the count rate from a sample of technetium-99 changes with time.



28

- (a) Alpha particles (α), beta particles (β) and gamma rays (γ) are types of nuclear radiation.

- (i) Which of the three types of radiation is the most strongly ionising?

.....


(1)

- (ii) What effect does nuclear radiation have on living cells?

.....

(1)

- (b) The diagrams show a G-M tube and counter used to measure the radiation emitted from a source. Both diagrams show the reading on the counter one minute after it was switched on.



Explain why the counter readings show that the source is giving out only gamma radiation.

.....

.....

.....

.....

(2)

- (c) The box gives information about the radioactive isotope technetium-99.

| |
|--|
| <p>Type of radiation emitted: gamma</p> <p><i>Half-life:</i> 6 hours</p> <p>Used as a medical tracer</p> |
|--|

What is meant by the term *half-life*?

.....

.....

(1)

- (d) To study the blood flow in a patient's lungs, a doctor injects a small quantity of a technetium-99 compound into the patient. The radiation emitted by the technetium-99 atoms is detected outside the patient's body.

Explain why a doctor would not use a radioactive isotope with a very short half-life, such as 2 seconds, as a medical tracer.

.....
.....
.....
.....

(2)
(Total 7 marks)

29

Some types of food are treated with *gamma* radiation. Low doses of radiation slow down the ripening of fresh fruit and vegetables while higher doses of radiation kill the bacteria that make the food go off.

- (a) (i) What is *gamma* radiation?

.....

(1)

- (ii) Food packed in crates or boxes can be treated using this method.

Why must a source that emits *gamma* radiation be used?

.....
.....

(1)

- (iii) A suitable source of gamma radiation is the isotope caesium 137.


Complete the following sentence by choosing the correct word from the box.

| | | |
|------------------|-----------------|----------------|
| electrons | neutrons | protons |
|------------------|-----------------|----------------|

An atom of caesium 137 has two more than an atom of caesium 135.

(1)

- (b) The diagram shows how a conveyor belt can be used to move food past the radioactive source.



- (i) How do the concrete walls reduce the radiation hazard to workers outside the food treatment area?

.....

(1)

- (ii) Suggest **one** way that the dose of radiation received by the food could be increased other than by changing the radioactive source.

.....

(1)

- (c) Some people may not like the idea of eating food treated with radiation.

- (i) What evidence could a food scientist produce to show that food treated with radiation is safe to eat?

.....

(2)

- (ii) The diagram shows the sign displayed on food treated with radiation.




Why is it important for people to know which foods have been treated with radiation?

.....
.....

(1)
(Total 8 marks)

30

The diagram shows a radiation detector and counter being used to measure background radiation. The number shows the count ten minutes after the counter was reset to zero.



- (i) Name **one** source of background radiation.

.....

(1)

- (ii) Calculate the average background radiation level, in counts per second. Show clearly how you work out your answer.

.....
.....

Background radiation level = counts per second

(2)
(Total 3 marks)

31

- (a) The table gives information about six radioactive isotopes.

| Isotope | Type of radiation emitted | Half-life |
|---------------|---------------------------|--------------------|
| hydrogen-3 | beta particle | 12 years |
| iridium-192 | gamma ray | 74 days |
| polonium-210 | alpha particle | 138 days |
| polonium-213 | alpha particle | less than 1 second |
| technetium-99 | gamma ray | 6 days |
| uranium-239 | beta particle | 24 minutes |

- (i) What is an alpha particle?

.....

(1)

- (ii) Two isotopes of polonium are given in the table. How do the nuclei of these two isotopes differ?

.....

(1)

- (iii) A doctor needs to monitor the blood flow through a patient's heart. The doctor injects a radioactive isotope into the patient's bloodstream. The radiation emitted by the isotope is then detected outside the body.

Which **one** of the isotopes in the table would the doctor inject into the bloodstream?

.....

Explain the reasons for your choice.

.....


.....

.....

.....

(3)

- (b) Igneous rock contains uranium-238 which eventually changes to the stable isotope lead-206. The graph shows how the percentage of uranium-238 nuclei present in an igneous rock changes with time.



A rock sample is found to have seven atoms of uranium-238 for every three atoms of lead-206. Use the graph to estimate the age of the rock. Show clearly how you obtain your answer.


.....
.....

Age of rock = million years

(2)
(Total 7 marks)

32

The diagram shows a badge used to monitor radiation. It measures the amount of radiation a worker has been exposed to in one month.



(i) What is used inside the badge to detect radiation?

.....

(1)

(ii) What would indicate that the worker has been exposed to a high level of radiation as opposed to a low level of radiation?

.....

.....

(1)

(iii) Why is it important to monitor the amount of radiation the worker has been exposed to?


.....

.....

(1)
(Total 3 marks)

33

Radiation is around us all of the time. The pie chart shows the sources of this radiation.



(i) What is the main source of this radiation?

.....

(1)

(ii) What name is given to the radiation that is around us all of the time?

.....

(1)

(Total 2 marks)


34

The table shows how the count rate from a radioactive substance changes in 10 days.

| | | | | | | |
|---------------------------------|-----|-----|-----|-----|-----|----|
| Time in days | 0 | 2 | 4 | 6 | 8 | 10 |
| Count rate in counts per minute | 880 | 555 | 350 | 220 | 140 | 90 |

(a) Draw a graph of count rate against time.

The first two points have been plotted for you.



(3)

(b) (i) Use your graph to find out how long it takes for the count rate to fall from 880 counts per minute to 440 counts per minute.

Time = days

(1)

(ii) What is the half-life of this substance?

Half-life = days

(1)

- (c) The table gives the half-life and type of radiation given out by four different radioactive isotopes.

| Radioactive isotope | Half-life in days | Radiation given out |
|---------------------|-------------------|---------------------|
| bismuth-210 | 5.0 | beta |
| polonium-210 | 138.0 | alpha and gamma |
| radon-222 | 3.8 | alpha |
| thorium-234 | 24.1 | beta and gamma |

Some samples of each isotope have the same count rate today. Which sample will have the lowest count rate one month from today?

.....


Give a reason for your answer.

.....

(2)
(Total 7 marks)

35

The different sources of radiation to which we are exposed are shown in the pie chart. Some sources are natural and some artificial.



(i) Name **one** natural source of radiation shown in the pie chart.

.....

(1)

(ii) Name **one** artificial source of radiation shown in the pie chart.

.....

(1)


(Total 2 marks)

36

(a) A radioactive source can give out three types of emission:

- alpha particles
- beta particles
- gamma radiation.

The diagram shows the paths taken by the radiation emitted by two sources, **X** and **Y**.




What types of radiation are emitted by each of the sources?

Source **X** emits

Source **Y** emits

(2)

- (b) The diagram shows a disposable syringe sealed inside a plastic bag. After the bag has been sealed the syringe is sterilised using radiation.



Explain why radiation can be used to sterilise the syringe.

.....

.....


.....

.....

(3)
(Total 5 marks)

37

The pie chart shows the main sources of *background radiation*. Each source contributes to the average yearly radiation dose.



- (i) What is meant by the term *background radiation*?

.....

.....

(1)



(ii) Suggest why an airline pilot is likely to get a higher than average yearly radiation dose.

.....
.....
.....

(2)
(Total 3 marks)

38

(a) The diagram shows three different boxes and three radioactive sources. Each source is stored in a different box.



Gamma source



Beta source




Alpha source

Draw lines to show which source should be stored in each box so that the risk of radiation leakage is a minimum.

(2)

- (b) A leak in an underground oil pipe can be found by injecting a radioactive isotope into the oil. The ground is then tested with a radiation detector and counter.



- (i) State the type of detector used.


.....

(1)

- (ii) Complete the sketch graph to show how the reading on the detector will change as it passes along the ground above the pipe.

(1)

- (c) Gamma radiation can be used to kill cancer cells inside a person's head. During the treatment the patient is kept perfectly still while the source of gamma radiation moves in a circle.



- (i) Why is a source of gamma radiation the most suitable for this treatment?

.....

(1)

- (ii) Suggest why a moving source of radiation is used rather than one which is kept stationary.

.....
.....
.....
.....

(2)

- (iii) Gamma radiation is an electromagnetic wave. Give **two** properties common to all electromagnetic waves.

1

.....

2

.....

(2)

(Total 9 marks)

39

- (a) The table shows the half-life of some *radioactive* isotopes.

| Radioactive isotope | Half-life |
|---------------------|------------|
| magnesium-27 | 10 minutes |
| sodium-24 | 15 hours |
| sulphur-35 | 87 days |
| cobalt-60 | 5 years |

- (i) What is meant by the term *radioactive*?

.....
.....

(1)

- (ii) Which **one** of the isotopes in the table could form part of a compound to be used as a tracer in medicine? Explain the reason for your choice.

.....

.....


.....

.....

.....

(3)

- (iii) Draw a graph to show how the number of radioactive atoms present in the isotope cobalt-60 will change with time.



(3)

- (b) Nuclear power stations provide about 17% of the world's electricity. They add less than 1% to the total background levels of radiation. Some people are opposed to the use of nuclear fuels for the generation of electricity. Explain why.

.....

.....

.....

.....


.....

.....

(3)
(Total 10 marks)

40

- (a) The pie-chart shows the main sources of background radiation.



- (i) Which source in the pie-chart adds the smallest amount of radiation to background levels?

.....

(1)


- (ii) Name **two** natural sources of background radiation in the pie-chart.

1.

2.

(2)

- (b) The diagrams show how a radiation detector and counter can be used to measure radiation levels. In each case the numbers show the count one minute after the counter is switched on.



- (i) How many counts are just from background radiation?

.....

- (ii) How many counts are just from the source?

.....

- (iii) What type of radiation did the source give out?

.....

Give a reason for your answer.

.....

.....


.....

.....

(4)
(Total 7 marks)

41

- (a) A radiation detector and counter were used to detect and measure the radiation emitted from a weak source. The graph shows how the number of counts recorded in one minute changed with time.



- (i) Even though the readings from the counter were accurately recorded, not all the points fit the smooth curve. What does this tell us about the process of radioactive decay?

.....

(1)

- (ii) After ten minutes the number of counts recorded each minute is almost constant. Explain why.

.....

.....

.....

(2)

- (b) The radioactive isotope sodium-24 injected into the bloodstream can be used to trace blood flow to the heart. Sodium-24 emits both *beta particles* and *gamma rays*.

- (i) What is a *beta particle*?

.....

(1)

- (ii) What is a *gamma ray*?

.....

.....

(1)

- (iii) The count rate from a solution containing sodium-24 decreases from 584 counts per minute to 73 counts per minute in 45 hours. Calculate the half-life of sodium-24. Show clearly how you work out your answer.

.....

.....

.....

Half-life = hours

(3)

- (iv) Give **one** advantage of using sodium-24 to trace blood flow compared to using an isotope with a half-life of:

[A] ten years;

.....

(1)

[B] ten seconds.

.....


(1)

(Total 10 marks)

42

- (a) Tritium (${}^3_1\text{H}$) is an isotope of hydrogen. Tritium has a proton number of 1 and a mass number of 3.

- (i) The diagram below shows a simple model of a tritium atom. Complete the diagram by adding the names of the particles indicated by the labels.



(4)

- (ii) Explain how the nucleus of an ordinary hydrogen atom is different from the nucleus of a tritium atom. Ordinary hydrogen atoms (${}^1_1\text{H}$) have a mass number of 1.

.....

.....

.....

(2)


- (iii) Tritium is a radioactive substance which emits beta (β) radiation. Why do the atoms of some substances give out radiation?

.....

.....

(2)

- (b) Tritium is one of the elements found in the waste material of the nuclear power industry. The diagram below shows a worker behind a protective screen. The container holds a mixture of different waste materials which emit alpha (α), beta (β) and gamma (γ) radiation.



Suggest a suitable material for the protective screen. The material should prevent radiation from the container reaching the worker. Explain your answer.


.....

.....

.....

(2)
(Total 10 marks)

The chart below shows the sources of radiation in Britain.



(a) Give **two** sources of natural radioactivity from the chart.

.....

(2)

(b) How might the chart be used to reassure people that nuclear power is safe?

.....

(1)

(c) Some material is spilled on a bench. How could you find out if this material is radioactive?

.....

(2)

- (d) The table shows the proton number and mass number of two isotopes of iodine.

Iodine is found naturally in the world as the isotope I-127. Iodine-127 is not radioactive and is essential to life.

Other isotopes of iodine are formed in nuclear reactors. In the Chernobyl nuclear power station disaster in Ukraine an explosion caused a large quantity of the isotope iodine-131 to be released into the atmosphere. Iodine-131 is radioactive.

| | proton number | mass number |
|------------|---------------|-------------|
| iodine-127 | 53 | 127 |
| iodine-131 | 53 | 131 |

Explain, in terms of particles found in the nucleus, how an iodine-131 nucleus is different from an iodine-127 nucleus.

.....
.....

(2)

- (e) (i) Explain, as fully as you can, why iodine-131 could be harmful to our bodies.

.....
.....
.....
.....

(4)

- (ii) Iodine-131 and iodine-127 have the same chemical properties. Explain why this would be a problem if iodine-131 was taken into our bodies.

.....
.....
.....

(1)

(iii) The Chernobyl disaster took place in 1986. Do you think that iodine-131 from the disaster is still a threat to us today? Explain your answer.

.....
.....
.....

(3)
(Total 15 marks)

44

In some areas of the U.K. people are worried because their houses are built on rocks that release radon.

Read the information about radon.

- It is a gas.
- It is formed by the breakdown of radium.
- It emits alpha radiation.
- Each radon atom has 86 protons.
- Each radon atom has 136 neutrons.

Explain why it may be dangerous to live near rocks that release radon.

To gain full marks in this question you should write your ideas in good English. Put them into a sensible order and use the correct scientific words.

.....
.....
.....
.....

(Total 3 marks)

45

The picture shows a man at work in a factory that uses radioactive materials.



The radioactive material is kept behind glass shields. The man wears gloves so that he cannot touch the radioactive material directly.

Explain, as fully as you can, why these precautions are taken.

To gain full marks in this question you should write your ideas in good English. Put them into a sensible order and use the correct scientific words.

.....

.....

.....

.....

.....

.....

(Total 4 marks)

46

The table gives the properties of some radionuclides (radioactive isotopes).

| Radionuclide | Half life | Main type of radiation emitted |
|---------------|--------------|--------------------------------|
| Radon-220 | 54.5 seconds | Alpha |
| Americium-241 | 433 years | Alpha |
| Phosphorus-32 | 14 days | Beta |
| Strontium-90 | 28 years | Beta |
| Technetium-99 | 6 hours | Gamma |
| Cobalt-60 | 5 years | Gamma |

(i) Which radionuclide would be best for monitoring the thickness of aluminium foil?

.....

Explain the reason for your answer.

.....
.....
.....

(2)

(ii) Which radionuclide would be best for acting as a tracer inside the human body?

.....

Explain the reason for your answer.

.....
.....
.....

(2)


(Total 4 marks)

47

The isotope of sodium with a mass number of 24 is radioactive. The following data were obtained in an experiment to find the half-life of sodium-24.

| Time in hours | Count rate in counts per minute |
|---------------|---------------------------------|
| 0 | 1600 |
| 10 | 1000 |
| 20 | 600 |
| 30 | 400 |
| 40 | 300 |
| 50 | 150 |
| 60 | 100 |


- (a) Draw a graph of the results and find the half-life for the isotope. On the graph show how you obtain the half-life.



Half-life = hours

(4)

- (b) Sodium-24 decays by beta emission. The G.M. tube used in the experiment is shown in the diagram. Each beta particle which gets through the glass causes a tiny electric current to pass in the circuit connected to the counter.



- (i) Why must the glass wall of the G.M. tube be very thin?

.....

(1)

- (ii) Why is this type of arrangement of no use if the radioactive decay is by alpha emission?

.....

(1)

- (c) Sodium chloride solution is known as saline. It is the liquid used in 'drips' for seriously-ill patients. Radioactive sodium chloride, containing the isotope sodium-24, can be used as a tracer to follow the movement of sodium ions through living organisms.

Give **one** advantage of using a sodium isotope with a half-life of a few hours compared to using an isotope with a half-life of:

- (i) five years;

(1)

- (ii) five seconds.

(1)

(Total 8 marks)


48

Doctors sometimes need to know how much blood a patient has.

They can find out by using a radioactive solution.

After measuring how radioactive a small syringe-full of the solution is they inject it into the patient's blood.

YOUR BLOOD CIRCULATION



They then wait for 30 minutes so that the solution has time to become completely mixed into the blood.

Finally, they take a syringe-full of blood and measure how radioactive it is.

Example:

If the doctor injects 10 cm^3 of the radioactive solution and this is diluted 500 times by the blood there must be $10 \times 500 = 5000 \text{ cm}^3$ of blood.

(a) After allowing for background radiation:

- 10 cm³ of the radioactive solution gives a reading of 7350 counts per minute;
- a 10 cm³ sample of blood gives a reading of 15 counts per minute.

Calculate the volume of the patient's blood.
(Show your working.)


.....

.....

.....

.....

(4)



Radiation from radioactive substances can harm your body cells.

(b) The doctor's method of estimating blood volume will not be completely accurate. Write down **three** reasons for this.

- 1
- 2
- 3

(3)

(c) The doctors use a radioactive substance which loses half of its radioactivity every six hours. Explain why this is a suitable radioactive substance to use.

.....

.....

(2)
(Total 9 marks)

49

(a) Sam and Kris are arguing about alpha and gamma radiation.

Sam says that alpha radiation is more dangerous.

Kris disagrees. He thinks that gamma radiation is more dangerous. What do you think? Explain your answer as fully as you can.

.....
.....
.....
.....
.....
.....

(4)

(b) Cancer cells in a particular organ of the body can be killed by injecting a radioactive substance which is absorbed by that organ.

What other features must the radioactive substance have to make it suitable for this job?

.....
.....

(2)

(c) Radon is a radioactive gas with a half-life of 3.6 days. It often seeps into buildings from the ground.

Estimate how long it takes for 99% of a sample of radon gas to decay. (Show your working.)

.....
.....
.....
.....
.....

(2)

(Total 8 marks)