

Answer ALL questions. Write your answers in the spaces provided.

1. [In this question position vectors are given relative to a fixed origin O]

At time t seconds, where $t \geq 0$, a particle, P , moves so that its velocity $\mathbf{v} \text{ m s}^{-1}$ is given by

$$\mathbf{v} = 6t\mathbf{i} - 5t^{\frac{3}{2}}\mathbf{j}$$

When $t = 0$, the position vector of P is $(-20\mathbf{i} + 20\mathbf{j}) \text{ m}$.

- (a) Find the acceleration of P when $t = 4$

(3)

- (b) Find the position vector of P when $t = 4$

(3)

$$a) \quad \underline{\underline{a}} = \frac{d\mathbf{v}}{dt} = 6\mathbf{i} - 5\left(\frac{3}{2}\right)t^{\frac{1}{2}}\mathbf{j}$$

$$= 6\mathbf{i} - \frac{15}{2}t^{\frac{1}{2}}\mathbf{j}$$

$$t=4 \Rightarrow \underline{\underline{a}} = 6\mathbf{i} - 15\mathbf{j}$$

$$b) \quad \underline{\underline{s}} = \int \underline{\underline{v}} dt$$

$$= \int 6t\mathbf{i} - 5t^{\frac{3}{2}}\mathbf{j} dt$$

$$= 3t^2\mathbf{i} - 2t^{\frac{5}{2}}\mathbf{j} + \underline{\underline{s}}_0$$

$$t=0:$$

$$-20\mathbf{i} + 20\mathbf{j} = \underline{\underline{s}}_0$$

$$\Rightarrow \underline{\underline{s}} = (3t^2 - 20)\mathbf{i} + (20 - 2t^{\frac{5}{2}})\mathbf{j}$$

$$t=4:$$

$$\underline{\underline{s}} = (48 - 20)\mathbf{i} + (20 - 64)\mathbf{j} \\ = 28\mathbf{i} - 44\mathbf{j}$$



2. A particle, P , moves with constant acceleration $(2\mathbf{i} - 3\mathbf{j})\text{ms}^{-2}$

At time $t = 0$, the particle is at the point A and is moving with velocity $(-\mathbf{i} + 4\mathbf{j})\text{ms}^{-1}$

At time $t = T$ seconds, P is moving in the direction of vector $(3\mathbf{i} - 4\mathbf{j})$

(a) Find the value of T .

(4)

At time $t = 4$ seconds, P is at the point B .

(b) Find the distance AB .

(4)

a) $\underline{v} = \underline{u} + \underline{a}t.$

$$\underline{v} = (-\underline{i} + 4\underline{j}) + (2\underline{i} - 3\underline{j})T.$$

$$\frac{4 - 3T}{-1 + 2T} = \frac{-4}{3}.$$

$$12 - 9T = 4 - 8T \\ \Rightarrow T = 8.$$

b) $\underline{s} = \underline{u}t + \frac{1}{2}\underline{a}t^2.$

$$\underline{s}_{AB} = (-\underline{i} + 4\underline{j})t + (\underline{i} - \frac{3}{2}\underline{j})t^2$$

$$t = 4:$$

$$\underline{s}_{AB} = -4\underline{i} + 16\underline{j} + 16\underline{i} - 24\underline{j} \\ = 12\underline{i} - 8\underline{j}.$$

$$\Rightarrow |AB| = \sqrt{12^2 + 8^2} = \sqrt{208} = 14.42\text{m}.$$



3.




Figure 1

Two blocks, A and B , of masses $2m$ and $3m$ respectively, are attached to the ends of a light string.

Initially A is held at rest on a fixed rough plane.

The plane is inclined at angle α to the horizontal ground, where $\tan \alpha = \frac{5}{12}$

The string passes over a small smooth pulley, P , fixed at the top of the plane.

The part of the string from A to P is parallel to a line of greatest slope of the plane. Block B hangs freely below P , as shown in Figure 1.

The coefficient of friction between A and the plane is $\frac{2}{3}$

The blocks are released from rest with the string taut and A moves up the plane.

The tension in the string immediately after the blocks are released is T .

The blocks are modelled as particles and the string is modelled as being inextensible.

(a) Show that $T = \frac{12mg}{5}$ (8)

After B reaches the ground, A continues to move up the plane until it comes to rest before reaching P .

(b) Determine whether A will remain at rest, carefully justifying your answer. (2)

(c) Suggest two refinements to the model that would make it more realistic. (2)

$$a) \quad R_A = 2mg \cos \alpha = \frac{24}{13} mg.$$

F_{rA} is limiting.

$$F_{rA} = \frac{2}{3} \times \frac{24}{13} mg = \frac{48}{39} mg.$$



Question 3 continued

$$A: T - (2mg \sin \alpha + \frac{48}{39} mg) = 2ma.$$

$$\Rightarrow T - (\frac{30}{39} + \frac{48}{39})mg = 2ma$$

$$\Rightarrow T - 2mg = 2ma.$$

$$\Rightarrow 3T - 6mg = 6ma$$

$$B: 3mg - T = 3ma$$

$$\Rightarrow 6mg - 2T = 6ma.$$

$$\Rightarrow 6mg - 2T = 3T - 6mg$$

$$\Rightarrow 12mg = 5T$$

$$\Rightarrow T = \frac{12mg}{5}.$$

$$b) F_{\max} = \frac{16}{13} mg.$$

$$2mg \sin \alpha = \frac{10}{13} mg.$$

$$\frac{16}{13} mg > \frac{10}{13} mg,$$

so A will not move down the plane.

c) Allow the string to be extensible.
Involve friction at the pulley P.

4.




Figure 2

A ramp, AB , of length 8 m and mass 20 kg, rests in equilibrium with the end A on rough horizontal ground.

The ramp rests on a smooth solid cylindrical drum which is partly under the ground. The drum is fixed with its axis at the same horizontal level as A .

The point of contact between the ramp and the drum is C , where $AC = 5$ m, as shown in Figure 2.

The ramp is resting in a vertical plane which is perpendicular to the axis of the drum, at an angle θ to the horizontal, where $\tan \theta = \frac{7}{24}$.

The ramp is modelled as a uniform rod.

(a) Explain why the reaction from the drum on the ramp at point C acts in a direction which is perpendicular to the ramp. (1)

(b) Find the magnitude of the resultant force acting on the ramp at A . (9)

The ramp is still in equilibrium in the position shown in Figure 2 but the ramp is not now modelled as being uniform.

Given that the centre of mass of the ramp is assumed to be closer to A than to B ,

(c) state how this would affect the magnitude of the normal reaction between the ramp and the drum at C . (1)

a) The drum is smooth, so there is no friction at this point C .



Question 4 continued

- b) Resolving vertically: $R_A + R_C \cos \theta = 20g$.
Resolving horizontally: $F_r = R_C \sin \theta$

Moments about A:

$$20g \times 4 \cos \theta = 5 R_C.$$

$$\Rightarrow R_C = 16g \cos \theta = \frac{384}{25} g.
(= 150.528 \text{ N}).$$

$$R_A = 20g - R_C \cos \theta
= 51.49312 \text{ N}.$$

$$F_r = 150.528 \times \frac{7}{25} = 42.14784$$

$$\Rightarrow |\text{Force at A}| = \sqrt{51.49312^2 + 42.14784^2}
= \sqrt{4427.981824}
= 66.54 \text{ N}.$$

- c) The magnitude of the reaction at C, " R_C " would decrease.



5.




Figure 3

The points A and B lie 50 m apart on horizontal ground.

At time $t = 0$ two small balls, P and Q , are projected in the vertical plane containing AB .

Ball P is projected from A with speed 20 m s^{-1} at 30° to AB .

Ball Q is projected from B with speed $u \text{ m s}^{-1}$ at angle θ to BA , as shown in Figure 3.

At time $t = 2$ seconds, P and Q collide.

Until they collide, the balls are modelled as particles moving freely under gravity.

(a) Find the velocity of P at the instant before it collides with Q .

(6)

(b) Find

(i) the size of angle θ ,

(ii) the value of u .

(6)

(c) State one limitation of the model, other than air resistance, that could affect the accuracy of your answers.

(1)

a) A: Horizontal: $20 \cos 30^\circ$.
 Vertical: $u = 20 \sin 30^\circ$, $a = -9.8$, $t = 2$, $v = ?$
 $v = 20 \sin 30^\circ - 19.6 = -9.6$

$$|V_A| = \sqrt{(20 \cos 30^\circ)^2 + 9.6^2} = 19.8 \text{ m s}^{-1}$$

b) i) Horizontal: A: $S = S_A$, $u = 20 \cos 30^\circ$, $a = 0$, $t = 2$
 B: $S = S_B$, $u = u \cos \theta$, $a = 0$, $t = 2$.

$$S_A + S_B = 50.$$

$$S_A = 40 \cos 30^\circ$$

$$S_B = 2u \cos \theta.$$

$$\Rightarrow 2u \cos \theta = 50 - 20\sqrt{3}.$$

$$\Rightarrow u \cos \theta = 25 - 10\sqrt{3}.$$



Question 5 continued

Vertical: A: $s = s_{\Delta}$, $u = 20 \sin 30^\circ$, $a = -g$, $t = 2$.

B: $s = s$, $u = u \sin \theta$, $a = -g$, $t = 2$.

A: $s = 20 - 2g$

B: $s = 2u \sin \theta - 2g$.

$$\Rightarrow 2u \sin \theta = 20$$

$$\Rightarrow u \sin \theta = 10.$$

$$u \sin \theta = 10, \quad u \cos \theta = 25 - 10\sqrt{3}$$

$$\Rightarrow u^* = \sqrt{10^2 + (25 - 10\sqrt{3})^2} = 12.609 \text{ m s}^{-1}.$$

$$\tan \theta = \frac{10}{25 - 10\sqrt{3}}$$

$$\Rightarrow \theta = \tan^{-1}\left(\frac{10}{25 - 10\sqrt{3}}\right) = 52.478^\circ.$$

- c) The model doesn't consider the fact that the balls are not particles, so they do not move freely under gravity.