

Mark Scheme (Results)

Summer 2019

Pearson Edexcel International GCSE In Mathematics A (4MA1) Paper 2HR

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2019
Publications Code 4MA1_2HR_1906_MS
All the material in this publication is copyright
© Pearson Education Ltd 2019

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme.
 - Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Types of mark

- o M marks: method marks
- o A marks: accuracy marks
- o B marks: unconditional accuracy marks (independent of M marks)

Abbreviations

- o cao correct answer only
- ft follow through
- o isw ignore subsequent working
- o SC special case
- o oe or equivalent (and appropriate)
- o dep dependent
- o indep independent
- o eeoo each error or omission

No working

If no working is shown then correct answers normally score full marks If no working is shown then incorrect (even though nearly correct) answers score no marks.

· With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.

If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks.

Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks.

If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work. If there is a choice of methods shown, the method that gains the least marks should be marked.

If there is no answer on the answer line then check the working for an obvious answer.

Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. Incorrect cancelling of a fraction that would otherwise be correct.

It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect eg algebra.

Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

International GCSE Maths

Apart from Q2, 17, 18d, 20, & 24 (where the mark scheme states otherwise) the correct answer, unless obtained from an incorrect method, should be taken to imply a correct method

Que	estion	Working	Answer	Mark		Notes
1	(a)		x>-3	1	B1	Accept - 3 < x
	(b)	4 <i>y</i> − <i>y</i> ≤ 8 + 13	<i>y</i> ≤ 7 oe	2	M1	Arranging y 's on one side and the numbers on the other side. (allow $4y - y = 8 + 13$ oe or $4y - y < 8 + 13$ oe or $4y - y > 8 + 13$ oe or $4y - y \ge 8 + 13$ oe Allow $y \le 21/3$
						Total 3 mark

2	$\frac{17}{3}(-)\frac{11}{4}$ or $5\frac{8}{12}(-)2\frac{9}{12}$	3	M1	Sight of $\frac{17}{3}$ and $\frac{11}{4}$ or $5\frac{8}{12}$ and $2\frac{9}{12}$
	$\frac{68}{12} - \frac{33}{12}$ or $4\frac{20}{12} - 2\frac{9}{12}$		M1	or $\frac{68n}{12n} - \frac{33n}{12n}$
	$\frac{35}{12} = 2 \frac{11}{12}$			
	Alt: $(2.61)^{1/2}$		A1	Dep on M2
	$3 (+)(\frac{2}{3} - \frac{3}{4})$ $3 (+)(\frac{3}{12} - \frac{9}{12})$	 -	-·-M1	
	$3 (+) (\frac{3}{12} - \frac{9}{12}) 3 - \frac{1}{12} = 2 \frac{11}{12}$		M1	
	Alt: $4^{\frac{5}{2}}(-)2^{\frac{3}{2}}$		A1	Dep on M2
	Alt: $4\frac{5}{3}(-)2\frac{3}{4}$ $2(+)(\frac{5}{3}-\frac{3}{4})$ $2(+)(\frac{20}{12}-\frac{9}{12})$		M1	
	$= 2 \frac{11}{12}$	 		
			M1 A1	Dep on M2
				Total 3 marks

Que	stion	Working	Answer	Mark		Notes
3	(a)		-5, 5, 5, -5	2		rrect values
						2 then B1 for 2 or 3 correct values
	(b)			2		g at least 6 points correctly from their
					table d	ep on B1 in part(a)
			Fully correct curve			accept horizontal line at top of curve
					or strai	ght line segments
						Total 4 marks
4	(a)	40 ÷ 16 × 12 oe		2	M1 $40 \times \frac{12}{16}$	20
-	(α)	40 * 10 * 12 00			40 X 16	Je
			30		A1	
	(b)	525 ÷ 100 ²		2	M1 525	
	` ′				$\overline{100^2}$	
			0.0525 oe		A1 Accept	5.25 x 10 ⁻²
						Total 4 marks
	l		l			
5				2	M1 For (x +	a (x + b) where $ab = -36$ and a and b
					are inte	
			(x + 4) (x - 9)		A1 Ignore	extension to roots x = -4 & 9
						Total 2 marks
6	1	P(mint =) 1 - (0.35 + 0.32 + 0.12) {= 0.21}		3	M1 Or a co	rrect equation summing to 1
0		P(strawberry or mint =) 0.32 + "0.21"		5	M1 Dep	rrect equation summing to 1
		1 (Strawberry or minic =) 0.32 + 0.21	0.53 oe		A1 Allow 0	53/1
			0.55 00		Al Allow o	Total 3 marks
	ı					
7		55 ÷ (6 + 3 + 2) {= 5}		3	M1 Or $\frac{6}{11}$ X	55 (= 30) or $\frac{2}{11}$ x 55 (= 10)
		(6 x "E") (2 x "E")				
		(6 x "5") – (2 x "5")				for Won = 30 <u>and</u> Lost = 10
			20		_ ·	seen in a ratio 30 : 15: 10)
	ļ		20		A1	
						Total 3 marks

Que	stion	Working	Answer	Mark		Notes
8	(a)	$ \begin{array}{c c} A & & B \\ \hline 5 & \begin{pmatrix} 3^2 \\ 5^3 \\ 7 \end{pmatrix} 3^2 & 11 \end{array} $		2	M1	3 ² x 5 ³ x 7 oe or correct Venn diagram
			7875		A1	
	(b)	$ \begin{array}{c c} A & & B \\ \hline 5 & \begin{pmatrix} 3^2 \\ 5^3 \end{pmatrix} 3^2 & 11 \end{array} $		2	M1	3 ⁴ x 5 ⁴ x 7 x 11 oe or correct Venn diagram
			3 898 125		A1	
						Total 4 marks
			0.4.405			
9	(a)	6000000	8.4 × 10 ⁵	1	B1	
	(b)	60000000000000000000000000000000000000	7.5 × 10 ⁸	2	M1 A1	M1 for 60000000 or 0.08
			7.5 * 10		7(1	Total 3 marks
	I			l	1	
10		150000 x 0.82 ³	02705	3	M2	If not M2 then M1 for 1st year e.g 150000 x 0.82 (= 123000) or 150000 x 0.18 (= 27000) SC B1 for 150000 x 1.18 (= 177000) or 150000 x 1.18 ³ (= 246454.8)or 150000 x 0.54 (=81000) or 150000 x 0.46 (= 69000)
			82705		A1	Accept 82705.2 Total 3 marks
						<u> </u>
11		Gradient = (−)4 ÷ 2 oe		3	M1	Correct method to work out the gradient (\pm) accept $4 \div 2$ oe or " m " = 2
			y = -2x - 1 oe		A2	If not A2 then A1 for $L = -2x - 1$ or $-2x - 1$ or $y = 2x - 1$ or $y = -2x + c$
						Total 3 marks

Question	Working	Answer	Mark		Notes
12	$\sin 32 = \frac{BD}{3.1} \text{ oe}$		5	M1	A correct calculation involving BD
	$(BD =)3.1 \times \sin 32 (= 1.6427)$			M1	Accept 1.6 or better
	$\cos 42 = \frac{\text{"3.1 sin 32"}}{AB} \text{ oe or } \frac{AB}{\sin 90} = \frac{\text{"3.1 sin 32"}}{\sin 48} \text{ oe}$			M1	Dep or (AD =) "1.6 x tan 42 {= 1.479}
	$AB = \frac{\text{"3.1}\sin 32\text{"}}{\cos 42} \text{ or } AB = \frac{\text{"3.1}\sin 32\text{"}}{\sin 48}$			M1	Or (AB =) $\sqrt{1.479^2 + 1.6427^2}$
		2.21		A1	2.21053 (Accept 2.2 → 2.22)
					Total 5 marks
13 (a)	Plotting points from table at ends of interval		2	M1	$\pm \frac{1}{2}$ sq (at least 5 points plotted correctly) Or a

13	(a)	Plotting points from table at ends of interval (40, 6), (50, 20), (60, 56), (70, 84), (80,95), (90, 100)		2	M1	$\pm^{1}/_{2}$ sq (at least 5 points plotted correctly) Or <u>all</u> points plotted consistently within each interval at the correct heights Accept cf graph which is not joined to (30,0)
		Points joined with curve or line segments	Correct cf diagram		A1	
	(b)	Use of graph at 50		2	M1	Use of graph at 50 walkers
			58 – 59		A1	No working shown and answer is within 58 – 59 award M1A1
	(c)	86 or 87 or 88 indicated on graph or stated		3	M1	Use of their graph at 72 minutes
		100 – "86" or 100 – "87" or 100 – "88"			M1	Dep e.g. 12, 13 or 14 walkers
			$\frac{12}{100}$ oe $\frac{13}{100}$ oe $\frac{14}{100}$		A1	0.12 → 0.14 inc, oe
						Total 7 marks

14	(a)	x ⁹ y ⁶	<i>x</i> ⁹ <i>y</i> ⁶	2	B1B1	Allow B1 if $(x^3y^2)^3$ or $(x^{36}y^{24})^{0.25}$ seen on answer
						line
	(b)	$3^n = \frac{3^n}{3^n}$		2	M1	for a correct first step e.g. 3 ^{2y} or 3 ^{-2y}
		3 ² y	n = x - 2y		A1	
						Total 4 marks

Question	Working	Answer	Mark		Notes
15	$ABD = 98^{\circ} \div 2 (= 49^{\circ}) \text{ or } ABC = 90^{\circ}$		4	M1	Correct angle stated or seen on diagram
	Angle at centre / middle is twice angle at circumference			B1	Dep M1
	Angle in a semicircle / from a diameter is 90° / right angle			B1	Dep M1
	DBC = (90 - 49) = 41	41°		A1	Correct answer + no reasons = M1A1
	- Alt:			.	
	180 – 98 (= 82°)				
	$OAD = 82 \div 2 (= 41^{\circ})$			M1	Correct angle stated or seen on diagram
	Base / bottom angles in an <u>isosceles triangle</u> are equal			B1	Dep M1
	DBC = 41°				
	Angles in the same segment / from the same chord (DC) are equal			B1	Dep M1
		41°		A1	Correct answer + no reasons = M1A1
	Alt:				
	- - DOC'= 180'-98 (= 82°) · · · · · · · · · · · · · · · · · · ·	+		M1	Correct angle stated or seen on diagram
	Angles on a straight line = 180°			B1	Dep M1
	DBC = 41°				
	Angle at centre / middle is twice angle at circumference			B1	Dep M1
		41°		A1	Correct answer + no reasons = M1A1
					Total 4 marks

16	(a)	$y = \frac{k}{x^2}$ condone proportion symbol in place of =		3	M1	Setting up a correct equation "k" ≠ 1
		$16 = \frac{k}{1.5^2}$ or $9 = \frac{k}{2^2}$ or $4 = \frac{k}{3^2}$ or $2.25 = \frac{k}{4^2}$			M1	Using the values from the table to find the value of the constant or " k " = 36
			$y = \frac{36}{r^2}$		A1	$\frac{36}{y^2}$ = M2 A0
	(b)	$x^2 = \frac{36}{144} \text{ or } x = \sqrt{\left(\frac{36}{144}\right)}$			M1	Substituting $y = 144$ into the correct equation and making x^2 or x the subject.
			0.5 oe	2	A1	cao
						Total 5 marks

Question	Working	Answer	Mark		Notes
17	(Term $n = \frac{1}{2}n(n+1)$ or		4		Algebraic representation of one of the two
	(Term $n+1=1$) $\frac{1}{2}(n+1)(n+2)$			M1	consecutive terms in sequence
	$\frac{1}{2}n(n+1) + \frac{1}{2}(n+1)(n+2)$ $\frac{1}{2}(n+1)(n+n+2) = \frac{1}{2}(n+1)(2n+2) \text{ or}$			M1	Adding two consecutive terms
	$\frac{1}{2}n^2 + \frac{1}{2}n + \frac{1}{2}n^2 + \frac{1}{2}n + n + 1 \to \frac{n^2 + 2n + 1}{2}$				Factorisation or multiplying out correctly to get to
				M1	$n^2 + 2n + 1$
		$(n+1)^2$ shown		A1	Dep on M3
					Total 4 marks

18	(a)		$\frac{3}{4}$ oe	1	B1	
	(b)	$\frac{x-5}{4(x-5)-3}$		2	M1	
			$\frac{x-5}{4x-23}$		A1	cao
	(c)	$y = \frac{x}{4x-3} \text{ or } x = \frac{y}{4y-3}$ $y(4x-3) = x \qquad \text{or} \qquad x(4y-3) = y$ $4xy - 3y = x \qquad \text{or} \qquad 4xy - 3x = y$ $4xy - x = 3y \qquad \text{or} \qquad 4xy - y = 3x$ $x(4y-1) = 3y \qquad \text{or} \qquad y(4x-1) = 3x$	$\frac{3x}{4x-1}$ oe	3	M1 M1 A1	Moving the denominator to the other side of the equation Factorising the variable on one side in a correct expression
	(d)	Tangent drawn at <i>x</i> = – 0.5 (G =) 18 ÷ 3 oe	5 → 7	3	M1 M1 A1	Drawing a tangent at $x = -0.5$ Correct method to work out the gradient of the tangent at $x = -0.5$ or $x = +0.5$ Dep on 1 st M1 SC B1 B1 for drawing a tangent at $x = +0.5$ and gradient $= -3 \rightarrow -4$
						Total 9 marks

Question	Working	Answer	Mark		Notes
19	$\frac{25}{2}\pi = \pi r^2 \times \frac{80}{360}$		6	M1	Equation of sector equal to $\frac{25 \pi}{2}$ or a
	2 360				calculation that leads to r or r^2
	r = 7.5				
	1 - 7.5			A1	
	$(APB =) 2 \times \pi \times .7.5" \times \frac{80}{360} (= 10.471)$	+	.	NAA 64	Dep on 1 st M1
	(APB =) 10.471 (=10 π /3)			M1 ft	Accept 10.5 or better
	$(AB^2) = "7.5"^2 + "7.5"^2 - (2 \times "7.5" \times "7.5" \times \cos 80)$.	M1ft	Dep on 1 st M1
	$ \operatorname{Or} \frac{AB}{\sin 80} = \frac{-^{\circ}7.5^{\circ}}{\sin 80} = \frac{\sin 50}{\sin 80} $				Correct equation to find AB (= 9.6) or AB^2 (=
	or (AB =) 2 x "7.5" x sin 40				93 or better) must use a clearly identified
	(AB =) 9.6418				radius value
	"9.6418" + "10.4719"			N 4 4 5 4	
				M1ft	 - Dep on 2 nd and 3 rd method marks
		20.f		AI	awrt 20.1
					Total 6 marks
20			3		•
	3.455 or 3.465 or 6.25 or 6.35			M1	Accept 3.4649 for 3.465 or 6.349 for
					6.35
	_6×3.465			M1	$\frac{6 \times \text{UB}_a}{\text{LP}_a \text{ UP}}$ where
	6.25 - 3.465			IVII	$LB_b - UB_a$ where
					$3.46 < UB_a \le 3.465$ and
					$6.25 \le LB_b < 6.3$
		7.46		A1	Dep M2 Accept 7.46499
					Total 3 marks
l			1	1	I
21	(LSF =) $\sqrt{240 \div 540}$ or $\frac{2}{3}$ or $\frac{3}{2}$ or 1.5 or 3 : 2 or 2 : 3		3	M1	
	$(\frac{2}{3})^3$ x 2025 oe accept 0.0.66 or better for 2/3				Full method leading to correct answer
	(3)	600		M1	
		600		AT	Total 2 mayles
		600		A1	Total 3 mar

Question		Working	Answer	Mark	Notes	
22		$-2(x^2 - 6x) + 5 \text{ or } -2(x^2 - 6x - 2.5)$		4	M1 Factorising by extracting – 2 in a correct expression	
		$-2[(x-3)^2 - 9 - 2.5]$ or $-2[(x-3)^2 - 9] + 5$			M1 Correct expression equivalent to $5 + 12x - 2x^2$	
		$-2[(x-3)^2-11.5]$ or $-2(x-3)^2+18+5$			M1 Correct expression equivalent to $5 + 12x - 2x^2$	
			$23 - 2(x - 3)^2$		Award full marks if a, b, and c are correctly stated and $23 - 2(x - 3)^2$ is not stated anywhere. SC B3 for $23 - 2(3 - x)^2$ SC B2 for $-2(x - 3)^2 + \text{constant}$ or $23 - 2(x + \text{constant})^2$ SC B1 for $-2(x + 3)^2 + \text{constant}$	
		Alt: _a + b(x² + 2cx + c²)			M1 Multiplying out expression correctly	
		$2bc = 12 \text{ or } a + bc^2 = 5 \text{ or } b = -2$ $2 \times -2 \times c = 12 \text{ or } c = -3$			M1 Equating coefficients or stating value of b Method to calculate c	
		$a + -2 \times (-3)^2 = 5$ or $a = 23$ seen			M1 Method to calculate <i>a</i> M1 SC P3 for 23 2 262 272	
			$23 - 2(x - 3)^2$		SC B3 for $23 - 2(3 - x)^2$	
					Total 4 marks	

Question	Working	Answer	Mark		Notes
23	360 = (10 x 10) + 4 x 0.5 x 10 x "h" oe h = 13			M1 A1	Finding the perpendicular height of a triangular face
	$AC = \sqrt{13^2 + 5^2} = (13.93 \text{ or } \sqrt{194}) \text{ or}$ $AO = \sqrt{13^2 - 5^2} = (12) \text{ or}$ $OC = (\sqrt{10^2 + 10^2}) \div 2 = (7.07 \text{ or } 5\sqrt{2}) \text{ or}$ $EC \text{ (oe)} = \sqrt{10^2 + 10^2} = (14.14 \text{ or } 10\sqrt{2})$			M2	Finding the accurate length of two sides relevant to finding correct angle. M2 for two sides found or M1 for one side. 1dp rounded or truncated.
	$tan^{-1} \left(\frac{12}{7.07}\right) \text{ or } cos^{-1} \left(\frac{7.07}{13.93}\right) \text{ or } sin^{-1} \left(\frac{12}{13.93}\right)$ or $cos^{-1} \left(\frac{13.93^2 + 7.07^2 - 12^2}{2 \times 13.93 \times 7.07}\right)$ or $cos^{-1} \left(\frac{13.93^2 + 14.14^2 - 13.93}{2 \times 13.93 \times 14.14}\right)$			M1	A correct trigonometric expression to find correct angle Accept $\tan \theta = \left(\frac{12}{7.0}\right)$ etc
		59.5°		A1	Accept 59.4° – 59.7°
					Total 6 marks

Question	Working	Answer	Mark	Notes	
24	$\frac{x-4}{x} \times \frac{x-5}{x-1} = 0.7$		5	M2	If not M2 then M1 for either $\frac{x-4}{x}$ or $\frac{x-5}{x-1}$
	$3x^{2} - 83x + 200 (= 0) \text{ oe}$ $\frac{83 \pm \sqrt{83^{2} - (4 \times 3 \times 200)}}{2 \times 3} \text{ or } (3x - 8)(x - 25) (= 0)$			A1 M1	Rearrangement of their quadratic to the form $ax^2 + bx + c$ (= 0) 1st step in solving the correct 3 term quadratic
	or (x - 83/6) ² + 200/3 - 83 ² /36 (=0)				Accept 25 only (dep on M3 if using algebra)
		25		A1	
	Alt: y = yellow marbles				If not M2 then M1 for either $\frac{y}{y+4}$ or $\frac{y-1}{y+3}$
	$\frac{y-1}{y+4} \times \frac{y-1}{y+3} = \overline{0.7}$			M2	Rearrangement of their quadratic to the form $ay^2 + by + c = 0$
	$3y^2 - 59y - 84 (= 0)$ oe $59 \pm \sqrt{59^2 - (4 \times 3 \times - 84)}$			A1	1 st step in solving the correct 3 term quadratic
	$\frac{59\pm\sqrt{59^2-(4\times3\times-84)}}{2\times3} \text{ or } (3y+4)(y-21)$ or $(y-59/6)^2-84/3-59^2/36$ (=0)			M1	Accept 25 and (day on M2 if units also has)
	y = 21 21+4				Accept 25 only (dep on M3 if using algebra) Give full marks if $\frac{21}{25} \times \frac{20}{24} = 0.7$ seen and 1st M2
		25		A1	scored NB: SC B1 for completing 1 st step in solving incorrect 3 term quadratic
					Total 5 marks

			Total for Paper: 100 marks

