



Please write clearly in block capitals.

Centre number

--	--	--	--	--

Candidate number

--	--	--	--

Surname

MODEL SOLUTIONS

Forename(s)

Candidate signature

I declare this is my own work.

# A-level MATHEMATICS

Paper 3

Friday 12 June 2020

Afternoon

Time allowed: 2 hours

## Materials

- You must have the AQA Formulae for A-level Mathematics booklet.
- You should have a graphical or scientific calculator that meets the requirements of the specification.

## Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer each question in the space provided for that question. If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work that you do not want to be marked.

## Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 100.

## Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- You do not necessarily need to use all the space provided.

For Examiner's Use

Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	
14	
15	
16	
17	
18	
<b>TOTAL</b>	




JUN207357301

PG/June20E8



7357/3

# MME.

## A Level Products Revision Cards



## Predicted Papers



Available to buy separately or as a bundle

## Section A

Answer all questions in the spaces provided.

- 1 Given that

$$\int_0^{10} f(x) dx = 7$$

deduce the value of

$$\int_0^{10} (f(x) + 1) dx$$

Circle your answer.

[1 mark]

-3

7

8

17

- 2 Given that

$$6 \cos \theta + 8 \sin \theta \equiv R \cos(\theta + \alpha)$$

find the value of  $R$ .

Circle your answer.

[1 mark]

6

8

10





14



3 Determine which one of these graphs does not represent  $y$  as a function of  $x$ .

Tick (✓) one box.

[1 mark]



Do not write  
outside the  
box.

Turn over ►



$$4 \quad p(x) = 4x^3 - 15x^2 - 48x - 36$$

4 (a) Use the factor theorem to prove that  $x - 6$  is a factor of  $p(x)$ .

[2 marks]

If  $x - 6$  is a factor, then  $p(6) = 0$ .

$$p(6) = 4(6)^3 - 15(6)^2 - 48(6) - 36$$

$$= 864 - 540 - 288 - 36$$

$$= 0$$

4 (b) (i) Prove that the graph of  $y = p(x)$  intersects the  $x$ -axis at exactly one point.

[4 marks]

$$p(x) = (x-6)(4x^2+9x+6)$$

$$\left( \begin{aligned} &= 4x^3 + 9x^2 + 6x - 24x^2 - 54x - 36 \\ &= 4x^3 - 15x^2 - 48x - 36 \end{aligned} \right) \Rightarrow$$

$$\text{Let } 4x^2 + 9x + 6 = 0$$

$$\Rightarrow \text{discriminant} = 9^2 - 4 \times 4 \times 6 < 0$$

$\therefore$  has no real solutions, thus the only solution to  $p(x) = 0$  is  $x = 6$ , so

$y = p(x)$  only ~~touches~~ intersects the  $x$ -axis at one point,  $(6, 0)$ .



---

---

---

---

---

---

---

---

---

---

4 (b) (ii) State the coordinates of this point of intersection.

[1 mark]

$(6, 0)$

---

---

---

---

---

---

---

Turn over for the next question

Turn over ►





- 5 The number of radioactive atoms,  $N$ , in a sample of a sodium isotope after time  $t$  hours can be modelled by

$$N = N_0 e^{-kt}$$

where  $N_0$  is the initial number of radioactive atoms in the sample and  $k$  is a positive constant.

The model remains valid for large numbers of atoms.

- 5 (a) It takes 15.9 hours for half of the sodium atoms to decay.

Determine the number of days required for at least 90% of the number of atoms in the original sample to decay.

[5 marks]

$$\text{At } t = 15.9, N = \frac{N_0}{2}$$

$$\Rightarrow \frac{N_0}{2} = N_0 e^{-15.9k}$$

$$\Rightarrow \frac{1}{2} = e^{-15.9k}$$

$$\Rightarrow \ln \frac{1}{2} = -15.9k$$

$$\Rightarrow k = \frac{\ln \frac{1}{2}}{-15.9} = 0.0436$$

and

$$0.1 N_0 = N_0 e^{-0.0436t}$$

$$\Rightarrow 0.1 = e^{-0.0436t}$$

$$\Rightarrow \ln 0.1 = -0.0436t$$

$$\Rightarrow t = \frac{\ln 0.1}{-0.0436} = 52.8 \text{ hours} = 2.2 \text{ days.}$$



- 5 (b) Find the percentage of the atoms remaining after the first week.

Give your answer to two significant figures.

[2 marks]

$$N = N_0 e^{-0.0436 \times 24 \times 7}$$

$$\Rightarrow N = N_0 \times 0.00658 \dots$$

$$\% \text{ remaining} = 0.066 \%$$

- 5 (c) Explain why the model can only provide an estimate for the number of remaining atoms.

[1 mark]

The model is continuous but the  
number of atoms is discrete.

- 5 (d) Explain why the model is invalid in the long run.

[1 mark]

The model will eventually predict less  
than one atom remaining, but never  
zero for large values of  $t$ .

Turn over for the next question


Turn over ►






- 6 The graph of  $y = f(x)$  is shown below.

Do not write  
outside the  
box




- 6 (a) Sketch the graph of  $y = f(-x)$

[2 marks]



6 (b) Sketch the graph of  $y = 2f(x) - 4$ 


[2 marks]

6 (c) Sketch the graph of  $y = f'(x)$ 

$$f'(x) = \text{gradient} \quad [3 \text{ marks}]$$

$$= \frac{6+2}{2+2} = 2$$

(between  $-2 \leq x \leq 2$ )



Turn over for the next question

Turn over ▶



7 (a) Using  ${}^n C_r = \frac{n!}{r!(n-r)!}$  show that  ${}^n C_2 = \frac{n(n-1)}{2}$

[2 marks]

$$r = 2 \Rightarrow {}^n C_2 = \frac{n!}{2!(n-2)!}$$

$$\Rightarrow {}^n C_2 = \frac{n \times (n-1) \times (n-2)! \times \dots}{2 \times (n-2)! \times (n-2)! \times \dots}$$

$$= \frac{n(n-1)}{2} \quad \square$$

7 (b) (i) Show that the equation

$$2 \times {}^n C_4 = 51 \times {}^n C_2$$

simplifies to

$$n^2 - 5n - 300 = 0$$

[3 marks]

$$2 \times {}^n C_4 = 51 \times {}^n C_2$$

$$\Rightarrow \frac{2n!}{4!(n-4)!} = \frac{51n(n-1)}{2}$$

$$\Rightarrow \frac{2n(n-1)(n-2)(n-3)}{4!} = \frac{51n(n-1)}{2}$$

$$\Rightarrow \frac{(n-2)(n-3)}{6} = 51$$

$$\Rightarrow n^2 - 5n + 6 = 306$$

$$\Rightarrow n^2 - 5n - 300 = 0$$



---

---

---

---

---

---

7 (b) (ii) Hence, solve the equation

$$2 \times {}^n C_4 = 51 \times {}^n C_2$$

[2 marks]

$$n = \frac{5 \pm \sqrt{5^2 - 4 \times 1 \times -300}}{2}$$

$$\Rightarrow n = 20 \text{ or } n = -15$$

---

---

---

---

---

---

---

---

---

---

Turn over for the next question

Turn over ►



8 The sum to infinity of a geometric series is 96

The first term of the series is less than 30

The second term of the series is 18

8 (a) Find the first term and common ratio of the series.

[5 marks]

$$ar = 18 \text{ (2nd term)} \Rightarrow a = \frac{18}{r}$$

$$\frac{a}{1-r} = 96 \text{ (sum to } \infty)$$

$$\Rightarrow \frac{\left(\frac{18}{r}\right)}{1-r} = 96$$

$$\Rightarrow \frac{18}{r} = 96 - 96r$$

$$\Rightarrow 18 = 96r - 96r^2$$

$$\Rightarrow 16r^2 - 16r + 3 = 0$$

$$\Rightarrow r = \frac{3}{4} \text{ or } r = \frac{1}{4}$$

$$\Rightarrow a = 24 \text{ or } a = 72$$

As  $a < 30$ , the  $a = 24$  and  $r = 0.75$

8 (b) (i) Show that the  $n$ th term of the series,  $u_n$ , can be written as

$$u_n = \frac{3^n}{2^{2n-5}}$$

[4 marks]

$$u_n = 24 \times \left(\frac{3}{4}\right)^{n-1}$$

$$= 24 \times \frac{3^{n-1}}{4^{n-1}}$$

$$= 3 \times 2^3 \times \frac{3^{n-1}}{2^{2(n-1)}}$$

$$= \frac{3^n}{2^{2n-5}}$$



8 (b) (ii) Hence show that

$$\log_3 u_n = n(1 - 2\log_3 2) + 5\log_3 2$$

[3 marks]

Taking  $\log_3$  of both sides: (from (b)(i))

$$\begin{aligned} \Rightarrow \log_3 u_n &= \log_3 \left( \frac{3^n}{2^{2n-5}} \right) \\ &= \log_3(3^n) - \log_3(2^{2n-5}) \\ &= n - (2n-5)\log_3(2) \\ &= n + (5-2n)\log_3(2) \\ &= n + 5\log_3 2 - 2n\log_3 2 \\ &= n(1 - 2\log_3 2) + 5\log_3 2. \end{aligned}$$

Turn over ►





9 (a) For  $\cos \theta \neq 0$ , prove that

$$\operatorname{cosec} 2\theta = \frac{1}{\sin 2\theta} \quad \operatorname{cosec} 2\theta + \cot 2\theta = \cot \theta$$

[4 marks]

$$\cot 2\theta = \frac{\cos 2\theta}{\sin 2\theta}$$

$\Rightarrow \operatorname{cosec} 2\theta + \cot 2\theta$  can be re-written as:

$$\Rightarrow = \frac{1}{\sin 2\theta} + \frac{\cos 2\theta}{\sin 2\theta}$$

$$= \frac{1}{2\sin\theta\cos\theta} + \frac{\cos^2\theta - \sin^2\theta}{2\sin\theta\cos\theta}$$

$$= \frac{1 + \cos^2\theta - \sin^2\theta}{2\sin\theta\cos\theta}$$

$$= \frac{2\cos^2\theta}{2\sin\theta\cos\theta}$$

$$= \frac{\cos\theta}{\sin\theta} = \cot\theta \quad \square$$

9 (b) Explain why

$$\cot \theta \neq \operatorname{cosec} 2\theta + \cot 2\theta$$

when  $\cos \theta = 0$

[1 mark]

$\operatorname{cosec} 2\theta$  and  $\cot \theta$  are undefined when  
 $\cos \theta = 0$ , as when  $\cos \theta = 0$ ,  $\sin 2\theta = 0$ .




## Section B

Do not write  
outside the  
box

Answer all questions in the spaces provided.

- 10 The probabilities of events  $A$ ,  $B$  and  $C$  are related, as shown in the Venn diagram below.

Find the value of  $x$ .

Circle your answer.

[1 mark]

0.11                      0.46                      0.54                      0.89

- 11 The table below shows the temperature on Mount Everest on the first day of each month.

Month	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Temperature ( $^{\circ}\text{C}$ )	-17	-16	-14	-9	-2	2	6	5	-3	-4	-11	-18

Calculate the standard deviation of these temperatures.

Circle your answer.

[1 mark]

-6.75                      5.82                      8.24                      67.85




- 12 The box plot below summarises the CO<sub>2</sub> emissions, in g/km, for cars in the Large Data Set from the London and North West regions.

London



North West



- 12 (a) Using the box plot, give **one** comparison of central tendency and **one** comparison of spread for the two regions.

[2 marks]

Comparison of central tendency

Median in London is greater than in the North West.

Range in London is less than in the North West.

Comparison of spread



Do not write outside the box

12 (b) Jaspal, an environmental researcher, used all of the data in the Large Data Set to produce a statistical comparison of the CO<sub>2</sub> and CO emissions in regions of England.

Using your knowledge of the Large Data Set, give **two** reasons why his conclusions may be invalid.

[2 marks]

Not all makes of car are included in  
 the data base.

---

Not all English regions are included.

---



---



---



---



---



---



---

Turn over for the next question

Turn over ►



- 13 Diedre is a head teacher in a school which provides primary, secondary and sixth-form education.

There are 200 teachers in her school.

The number of teachers in each level of education along with their gender is shown in the table below.

	Primary	Secondary	Sixth-form
Male	9	24	23
Female	35	85	24

- 13 (a) A teacher is selected at random. Find the probability that:

- 13 (a) (i) the teacher is female

[1 mark]

$$\frac{18}{25}$$

- 13 (a) (ii) the teacher is **not** a sixth-form teacher.

[1 mark]

$$\frac{153}{200}$$

- 13 (b) Given that a randomly chosen teacher is male, find the probability that this teacher is **not** a primary teacher.

[2 marks]

$$\text{Total male teachers} = 56$$

$$\text{Not primary} = 47/56$$



- 13 (c) Diedre wants to select three different teachers at random to be part of a school project.

Calculate the probability that all three chosen are secondary teachers.

[2 marks]

$$\frac{109}{200} \times \frac{108}{199} \times \frac{107}{198} = 0.16 \text{ (2.d.p.)}$$

---

---

---

---

---

---

---

Turn over for the next question

Turn over ►





14

It is known that a hospital has a mean waiting time of 4 hours for its Accident and Emergency (A&E) patients.

After some new initiatives were introduced, a random sample of 12 patients from the hospital's A&E Department had the following waiting times, in hours.

4.25 3.90 4.15 3.95 4.20 4.15

5.00 3.85 4.25 4.05 3.80 3.95

Carry out a hypothesis test at the 10% significance level to investigate whether the mean waiting time at this hospital's A&E department has changed.

You may assume that the waiting times are normally distributed with standard deviation 0.8 hours.

[7 marks]

$X$  = waiting time in hours.

$$H_0: \mu = 4$$

$$H_1: \mu \neq 4$$

$$\bar{x} = \frac{\sum x}{12} = 4.125$$

$$\text{Test statistic} = \frac{4.125 - 4}{\frac{0.8}{\sqrt{12}}} = 0.541$$

$$\text{Critical value} = 1.65$$

Accept  $H_0$  as  $0.541 < 1.65$ .

Therefore, there is insufficient evidence to suggest that the mean waiting time the A&E has changed.



- 15 A political party is holding an election to choose a new leader.

A statistician within the party decides to sample 70 party members to find their opinions of the leadership candidates.

There are 4735 members under 30 years old and 8565 members 30 years old and over.

The statistician wants to use a sample of 70 party members in the survey.

He decides to use a random stratified sample.

- 15 (a) Calculate how many of each age group should be included in his sample.

[2 marks]

$$\text{Under 30: } \frac{4735}{4735+8565} \times 70 = 24.9 \approx 25$$

$$\text{30+Over } \frac{8565}{4735+8565} \times 70 = 45.1 \approx 45$$

25 under 30 and 45 30 and over.

- 15 (b) Explain how he could collect the random sample of members under 30 years old.

[3 marks]

Number each member under 30 from

1 to 4735.

Using a calculator or random number generator,  
generate random 4 digit numbers.

Finish the process when you have 25  
different numbers.



- 16 An educational expert found that the correlation coefficient between the hours of revision and the scores achieved by 25 students in their A-level exams was 0.379

Her data came from a bivariate normal distribution.

Carry out a hypothesis test at the 1% significance level to determine if there is a positive correlation between the hours of revision and the scores achieved by students in their A-level exams.

The critical value of the correlation coefficient is 0.4622

[4 marks]

$$H_0: \rho = 0, H_1: \rho > 0$$

$$0.379 < 0.4622$$

There is insufficient evidence to reject  $H_0$ , therefore we accept  $H_0$ , that there <sup>isn't</sup> positive correlation between hours revised and A-level scores.

Turn over for the next question

Turn over ►



- 17 The lifetime of Zapple smartphone batteries,  $X$  hours, is normally distributed with mean 8 hours and standard deviation 1.5 hours.

17 (a) (i) Find  $P(X \neq 8)$

[1 mark]

1

17 (a) (ii) Find  $P(6 < X < 10)$

[1 mark]

0.818

17 (b) Determine the lifetime exceeded by 90% of Zapple smartphone batteries.

[2 marks]

$$\frac{x - \mu}{1.5} = \frac{x - 8}{1.5}$$

$$\Rightarrow x = 6.08$$



- 17 (c) A different smartphone, Kaphone, has its battery's lifetime,  $Y$  hours, modelled by a normal distribution with mean 7 hours and standard deviation  $\sigma$ .

25% of randomly selected Kaphone batteries last less than 5 hours.

Find the value of  $\sigma$ , correct to three significant figures.

[4 marks]

$$Z = -0.6745$$

$$\frac{5-7}{\sigma} = -0.6745$$

$$\sigma = 2.97$$

Turn over for the next question

Turn over ►



18 (a) (iii) a box contains at least 20 shirts which do **not** have a fabric defect.

[3 marks]

$$X \sim B(30, 0.70)$$

$$P(X \geq 20) = 1 - P(X \leq 19)$$

$$= 1 - 0.2696$$

$$= 0.7304$$

Do not write  
outside the  
box

Question 18 continues on the next page

Turn over ►





- 18 (b) Tiana wants to investigate the proportion,  $p$ , of defective shirts with a fabric defect. She wishes to test the hypotheses

$$H_0 : p = 0.3$$

$$H_1 : p < 0.3$$

She takes a random sample of 60 shirts with a defect and finds that  $x$  of them have a fabric defect.

- 18 (b) (i) Using a 5% level of significance, find the critical region for  $x$ .

[5 marks]

$$X \sim B(60, 0.30)$$

$$P(X \leq 11) = 0.0295 < 0.05$$

$$P(X \leq 12) = 0.0568 > 0.05$$

Critical region is therefore  $x \leq 11$ .



- 18 Tiana is a quality controller in a clothes factory. She checks for four possible types of defects in shirts.

Of the shirts with defects, the proportion of each type of defect is as shown in the table below.

Type of defect	Colour	Fabric	Sewing	Sizing
Probability	0.25	0.30	0.40	0.05

Shirts with defects are packed in boxes of 30 at random.

- 18 (a) Find the probability that:

- 18 (a) (i) a box contains exactly 5 shirts with a colour defect

[2 marks]

$$X \sim B(30, 0.25)$$

$$P(X=5) = 0.1047$$

- 18 (a) (ii) a box contains fewer than 15 shirts with a sewing defect

[2 marks]

$$X \sim B(30, 0.40)$$

$$P(X < 15) = P(X \leq 14)$$

$$= 0.8246$$



18 (b) (ii) In her sample she finds 13 shirts with a fabric defect.

Complete the test stating her conclusion in context.

[2 marks]

$13 > 11$  so we can accept  $H_0$ .

so there is insufficient evidence to suggest  
that the proportion of shirts with a fabric  
defect has decreased.

END OF QUESTIONS

Do not write  
outside the  
box.

