

Please check the examination details below before entering your candidate information

Candidate surname

Other names

Centre Number

Candidate Number

|  |  |  |  |  |
|--|--|--|--|--|
|  |  |  |  |  |
|--|--|--|--|--|

|  |  |  |  |
|--|--|--|--|
|  |  |  |  |
|--|--|--|--|

## Pearson Edexcel Level 3 GCE

Time 1 hour 30 minutes

Paper  
reference

**9FM0/01**

### Further Mathematics

Advanced

**PAPER 1: Core Pure Mathematics 1**

**You must have:**

Mathematical Formulae and Statistical Tables (Green), calculator

Total Marks

**Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for algebraic manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.**

#### Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided  
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear.  
Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

#### Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 10 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets  
– *use this as a guide as to how much time to spend on each question.*

#### Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P71776A

©2022 Pearson Education Ltd.

Q:1/1/1/1/



Pearson

1.  $f(z) = z^3 + az + 52$  where  $a$  is a real constant

Given that  $2 - 3i$  is a root of the equation  $f(z) = 0$

(a) write down the other complex root. (1)

(b) Hence

(i) solve completely  $f(z) = 0$

(ii) determine the value of  $a$  (4)

(c) Show all the roots of the equation  $f(z) = 0$  on a single Argand diagram. (1)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





2.

**In this question you must show all stages of your working.**

**Solutions relying entirely on calculator technology are not acceptable.**

Determine the values of  $x$  for which

$$64 \cosh^4 x - 64 \cosh^2 x - 9 = 0$$

Give your answers in the form  $q \ln 2$  where  $q$  is rational and in simplest form.

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



**Question 2 continued**

Lined area for writing the answer to Question 2.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**(Total for Question 2 is 4 marks)**



P 7 1 7 7 6 A 0 5 3 2





4. (a) Use the method of differences to prove that for  $n > 2$

$$\sum_{r=2}^n \ln\left(\frac{r+1}{r-1}\right) \equiv \ln\left(\frac{n(n+1)}{2}\right)$$

(4)

- (b) Hence find the exact value of

$$\sum_{r=51}^{100} \ln\left(\frac{r+1}{r-1}\right)^{35}$$

Give your answer in the form  $a \ln\left(\frac{b}{c}\right)$  where  $a$ ,  $b$  and  $c$  are integers to be determined.

(3)







**Question 4 continued**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





5.

$$\mathbf{M} = \begin{pmatrix} a & 2 & -3 \\ 2 & 3 & 0 \\ 4 & a & 2 \end{pmatrix} \quad \text{where } a \text{ is a constant}$$

- (a) Show that  $\mathbf{M}$  is non-singular for all values of  $a$ . (2)
- (b) Determine, in terms of  $a$ ,  $\mathbf{M}^{-1}$  (4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





6. (a) Express as partial fractions

$$\frac{2x^2 + 3x + 6}{(x + 1)(x^2 + 4)}$$

(3)

(b) Hence, show that

$$\int_0^2 \frac{2x^2 + 3x + 6}{(x + 1)(x^2 + 4)} dx = \ln(a\sqrt{2}) + b\pi$$

where  $a$  and  $b$  are constants to be determined.

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





Question 6 continued

Ruled writing area for the answer to Question 6.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





**Question 6 continued**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Lined writing area for the answer to Question 6.

(Total for Question 6 is 7 marks)



P 7 1 7 7 6 A 0 1 7 3 2



Question 7 continued

Handwritten area with horizontal lines for writing.

DO NOT WRITE IN THIS AREA  
DO NOT WRITE IN THIS AREA  
DO NOT WRITE IN THIS AREA



Question 7 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 7 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 7 is 7 marks)



8. (a) Given

$$z^n + \frac{1}{z^n} = 2 \cos n\theta \quad n \in \mathbb{N}$$

show that

$$32 \cos^6 \theta \equiv \cos 6\theta + 6 \cos 4\theta + 15 \cos 2\theta + 10 \quad (5)$$




Figure 1




Figure 2

Figure 1 shows a solid paperweight with a flat base.

Figure 2 shows the curve with equation

$$y = H \cos^3 \left( \frac{x}{4} \right) \quad -4 \leq x \leq 4$$

where  $H$  is a positive constant and  $x$  is in radians.

The region  $R$ , shown shaded in Figure 2, is bounded by the curve, the line with equation  $x = -4$ , the line with equation  $x = 4$  and the  $x$ -axis.

The paperweight is modelled by the solid of revolution formed when  $R$  is rotated  $180^\circ$  about the  $x$ -axis.

Given that the maximum height of the paperweight is 2 cm,

- (b) write down the value of  $H$ . (1)
- (c) Using algebraic integration and the result in part (a), determine, in  $\text{cm}^3$ , the volume of the paperweight, according to the model. Give your answer to 2 decimal places.

[Solutions based entirely on calculator technology are not acceptable.] (5)

- (d) State a limitation of the model. (1)

---



---



---



---



---



Question 8 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Lined area for writing the answer to Question 8.



**Question 8 continued**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA







9. (i) (a) Explain why  $\int_0^{\infty} \cosh x \, dx$  is an improper integral. (1)

(b) Show that  $\int_0^{\infty} \cosh x \, dx$  is divergent. (3)

(ii)  $4 \sinh x = p \cosh x$  where  $p$  is a real constant

Given that this equation has real solutions, determine the range of possible values for  $p$  (2)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**Question 9 continued**

Lined writing area for the response to Question 9.

**(Total for Question 9 is 6 marks)**



10.




Figure 3

The motion of a pendulum, shown in Figure 3, is modelled by the differential equation

$$\frac{d^2\theta}{dt^2} + 9\theta = \frac{1}{2}\cos 3t$$

where  $\theta$  is the angle, in radians, that the pendulum makes with the downward vertical,  $t$  seconds after it begins to move.

(a) (i) Show that a particular solution of the differential equation is

$$\theta = \frac{1}{12}t \sin 3t \tag{4}$$

(ii) Hence, find the general solution of the differential equation. (4)

Initially, the pendulum

- makes an angle of  $\frac{\pi}{3}$  radians with the downward vertical
- is at rest

Given that, 10 seconds after it begins to move, the pendulum makes an angle of  $\alpha$  radians with the downward vertical,

(b) determine, according to the model, the value of  $\alpha$  to 3 significant figures. (4)

Given that the true value of  $\alpha$  is 0.62

(c) evaluate the model. (1)

The differential equation

$$\frac{d^2\theta}{dt^2} + 9\theta = \frac{1}{2}\cos 3t$$

models the motion of the pendulum as moving with forced harmonic motion.

(d) Refine the differential equation so that the motion of the pendulum is simple harmonic motion. (1)









**Question 10 continued**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 10 is 14 marks)

**TOTAL FOR PAPER IS 75 MARKS**

